



King's Research Portal

DOI:

[10.1097/j.pain.0000000000001858](https://doi.org/10.1097/j.pain.0000000000001858)

Document Version

Peer reviewed version

[Link to publication record in King's Research Portal](#)

Citation for published version (APA):

Hoffmann, J. R., Miller, S., Martins-Oliveira, M., Akerman, S., Suprongsinchai, W., Sun, H., Shi, L., Wang, J., Zhu, D., Lehto, S., Liu, H., Yin, R., Moyer, B. D., Xu, C., & Goadsby, P. (2020). PAC1 receptor blockade reduces central nociceptive activity: new approach for primary headache? *Pain*, 161(7), 1670-1681.
<https://doi.org/10.1097/j.pain.0000000000001858>

Citing this paper

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

General rights

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal

Take down policy

If you believe that this document breaches copyright please contact librarypure@kcl.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

PAIN

PAC1 receptor blockade reduces central nociceptive activity: new approach for primary headache? --Manuscript Draft--

Manuscript Number:	PAIN-D-19-01173R1
Full Title:	PAC1 receptor blockade reduces central nociceptive activity: new approach for primary headache?
Article Type:	Research Paper
Keywords:	Pituitary Adenylate Cyclase Activating Peptide Sumatriptan Headache Migraine Cluster headache Trigeminal Activation Monoclonal Antibody
Corresponding Author:	Peter J Goadsby, MD PhD King's College London London, UNITED KINGDOM
Corresponding Author Secondary Information:	
Corresponding Author's Institution:	King's College London
Corresponding Author's Secondary Institution:	
First Author:	Jan Hoffmann, MD, PhD
First Author Secondary Information:	
Order of Authors:	<div>Jan Hoffmann, MD, PhD</div> <div>Silke Miller, PhD</div> <div>Margarida Martins-Oliveira, PhD</div> <div>Simon Akerman, PhD</div> <div>Weera Suprongsinchai, PhD</div> <div>Hong Sun, MD</div> <div>Licheng Shi, PhD</div> <div>Judy Wang, PhD</div> <div>Dawn Zhu, PhD</div> <div>Sonya Lehto, PhD</div> <div>Hantao Liu, PhD</div> <div>Ruoyuan Yin, PhD</div> <div>Bryan D. Moyer, PhD</div> <div>Cen Xu, PhD</div> <div>Peter J. Goadsby, MD, PhD</div>
Additional Information:	
Question	Response
Have you posted this manuscript on a preprint server (e.g., arXiv.org, BioXriv,	No

Reviewer 1

Excellent manuscript. I have only some minor points.

Thank you for sharing this judgement.

1) Please use "migraine patient" instead of "migraineur"

The word migraineur is correct (OED); there is no pressing, globally accepted view to insert two words when there is one English word that can be used.

2) page 11, line 44: "Experiments were conducted all studies were conducted..." Please correct

Thanks for picking up this mistake. It has now been corrected.

3) Could you please state in which years and in which laboratory exactly the experiments were conducted

The generation of the antibody Ab181, in vitro characterization of Ab181, pharmacokinetic and pharmacodynamic analyses as well as the immunohistochemistry were all conducted between 2011 and 2013 in the Amgen laboratories in Thousand Oaks, CA, USA. The electrophysiology experiments were all conducted between 2012 and 2013 at the University of California San Francisco, CA, USA. We have now stated this in the manuscript.

4) page 24: I think it is too early to conclude that there is only a peripheral mode of action. Given the differences between the animal model and humans regarding blood brain barrier etc., it might be that in humans there is a central site of action. Also, it is not yet proven that high concentrations of Ab181 are needed for the main mechanism. It could also be that the main mechanism is located in the CNS and that small amounts of Ab181 are sufficient.

We completely agree with this opinion. We have now amended the sentence accordingly.

Reviewer #2

The authors describe the development of a rodent-specific PAC1 receptor antibody (Ab181) and its effect on nociceptive neuronal activity in the trigemino-cervical complex. By doing so, they show that AB181 inhibits stimulus-evoked activity in the trigemino-cervical complex but not spontaneous neuronal activity. Using immunohistochemistry, they demonstrate that the antibody binds in the trigeminal ganglion and SPG but not within the CNS. The authors conclude that a PAC1 receptor antibody could provide potential benefit in treatment of primary headache. This manuscript adds important findings to the field because the potential benefit of PACAP antagonism in primary headache prevention is an ongoing discussion. However, I have few points to be addressed:

Results:

Dose response studies at 48h and 3.25 h. How did the authors choose these time points?

The time points were chosen to confirm that the receptor occupancy of Ab181 has reached and stayed at the desired pharmacodynamic measures during the entire duration of the electrophysiology experiments. We have now specified this in the manuscript.

Effect of Ab181 on neuronal activity in TCC: N number is missing. How many in vivo experiments in how many rats were performed? This is important for the interpretation of the main findings. In the methods section under the subheading general surgical preparation we had already stated that for the electrophysiological experiments we have used 24 animals. In each animal we have only recorded the electrophysiological signal from one site. We have now specified this in the manuscript.

Discussion:

Page 21 line14: The authors state that their data "provide a plausible biology for a clinical effect of PAC1 blockade". I would tone down this statement. The demonstration of inhibition of stimulated activity in the trigemino-cervical complex in an experimental animal model that might be associated with the pathophysiology of primary headaches does not justify such a conclusion.

We have now modified the sentence accordingly.

The authors state that the antagonism at the PAC1 receptor may offer a new strategy for the preventive treatment of primary headache disorders. This "new strategy" is somehow contradicted by the fact that a clinical phase-2 trial with a monoclonal PAC1 receptor antibody has already been reported to be ineffective in the preventive treatment of migraine. Instead of presenting a "new strategy" this paper supports the hypotheses that, despite the current clinical evidence, antagonism of the PAC1 receptor might offer some therapeutically gain in primary headache disorders. Accordingly, the authors discuss that PAC1 antibodies might work rather in migraine patients with autonomic symptoms or in cluster headache. I would suggest restructuring this paragraph under this aspect. Additionally, please provide the citation of the abstract of the phase 2 AMG 301 study.

We are aware of the fact that the phase 2 trial with AMG301 was negative. However, in our view it is much too early to conclude that the mechanism is ineffective in the preventive treatment of migraine. The main reason is that AMG 301 had a much lower affinity for the PAC1 receptor compared to Ab181. Given the amount of clinical and preclinical evidence supporting significant role of PACAP in the pathophysiology of migraine, it seems more likely that the lack of efficacy was based more on the specifications of AMG301 than on the mechanism itself. On top of this we have to consider the lack of distinction between migraine with and without autonomic symptoms and the complete lack of data on trigeminal autonomic cephalalgias. We will have more insight on the subject when the trials with monoclonal PACAP antibodies will have concluded and the results become available. Taken together, as we believe that AMG301 is not a good example to conclude that the mechanism is ineffective in migraine and as we have no data on trigeminoautonomic cephalalgias such as cluster headache, we have written that targeting the PAC1 receptor may offer a new strategy for the preventive treatment of primary headache disorders. Finally, we would like to cite the AMG301 data but they have only been presented as an e-poster at the EHF meeting 2019 and are not included in the abstract book published in the Journal of Headache and Pain. Hence, as much as we would like to, we can't cite the abstract.

Minor points

Introduction, line 44: The authors state that infusion of VIP "less reliably" induces migraine-like attacks. VIP does not induce migraine attacks as also demonstrated by the cited paper by Rahmann et al, Cephalalgia 2008 ("None of the subjects reported a migraine attack after VIP infusion."). Please correct.

We are aware of the work of Rahmann et al. However, the data is not that clear since Amin et al (Brain 2014; 137: 779-794) have shown that VIP may trigger attacks in some individuals. We therefore chose to use the terms "less reliably".

Results: Fig 3.2. -> Please add in the results text the referral to the designated graph (A,B,C,D)

We have now corrected this.

1 Pituitary adenylate cyclase activating polypeptide-38 (PACAP38) may play an important role
2 in primary headaches. Preclinical evidence suggests that PACAP38 modulates trigeminal
3 nociceptive activity mainly through PAC₁ receptors while clinical studies report that plasma
4 concentrations of PACAP38 are elevated in spontaneous attacks of cluster headache and
5 migraine and normalize after treatment with sumatriptan. Intravenous infusion of PACAP38
6 induces migraine-like attacks in migraineurs and cluster-like attacks in cluster headache
7 patients.
8

9
10 A rodent-specific PAC₁ receptor antibody Ab181 was developed and its effect on nociceptive
11 neuronal activity in the trigeminocervical complex was investigated *in vivo* in an
12 electrophysiological model relevant to primary headaches.
13

14 Ab181 is potent and selective at the rat PAC₁ receptor and provides near maximum target
15 coverage at 10 mg/kg for more than 48 hours. Without affecting spontaneous neuronal
16 activity, Ab181 effectively inhibits stimulus-evoked activity in the trigeminocervical complex.
17

18 Immunohistochemical analysis revealed its binding in the trigeminal ganglion and
19 sphenopalatine ganglion but not within the CNS suggesting a peripheral site of action.
20

21 The pharmacological approach using a specific PAC₁ receptor antibody could provide a novel
22 mechanism with a potential clinical efficacy in the treatment of primary headaches.
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

PAC₁ receptor blockade reduces central nociceptive activity- new approach for primary headache?

Jan Hoffmann MD PhD^{1,2}, Silke Miller PhD³, Margarida Martins-Oliveira PhD²,
Simon Akerman PhD^{2,¶}, Weera Suprongsinchai PhD², Hong Sun MD³,
Licheng Shi PhD³, Judy Wang PhD³, Dawn Zhu PhD³, Sonya Lehto PhD³,
Hantao Liu PhD³, Ruoyuan Yin PhD³, Bryan D. Moyer PhD³,
Cen Xu PhD^{3*} and Peter J. Goadsby MD PhD^{1,2*}

¹Basic and Clinical Neuroscience, Institute of Psychiatry, Psychology and Neuroscience,
King's College London, UK

²UCSF Headache Group - Department of Neurology, University of California, San Francisco,
San Francisco, CA, USA

³Department of Neuroscience, Amgen Inc.,
Thousand Oaks, CA, USA

Current address

[¶]Department of Neural and Pain Sciences, University of Maryland Baltimore, Baltimore, MD,
USA

*** These senior authors contributed equally to this manuscript**

Characters- Title: 87; Running Head: 26

Words- Abstract 186; Introduction 487; Discussion 1412; Total 6050

Figures-7 (color- 2); Tables- 1

Number of pages: 46

Correspondence to:

Prof. Peter J. Goadsby
NIHR - Wellcome Trust Clinical Research Facility
King's College Hospital
London SE5 9PJ
UK
Email: peter.goadsby@kcl.ac.uk

PAC₁ receptor blockade reduces central nociceptive activity- new approach for primary headache?

Jan Hoffmann MD PhD^{1,2}, Silke Miller PhD³, Margarida Martins-Oliveira PhD²,
Simon Akerman PhD^{2,¶}, Weera Suprongsinchai PhD², Hong Sun MD³,
Licheng Shi PhD³, Judy Wang PhD³, Dawn Zhu PhD³, Sonya Lehto PhD³,
Hantao Liu PhD³, Ruoyuan Yin PhD³, Bryan D. Moyer PhD³,
Cen Xu PhD^{3*} and Peter J. Goadsby MD PhD^{1,2*}

¹Basic and Clinical Neuroscience, Institute of Psychiatry, Psychology and Neuroscience,
King's College London, UK

²UCSF Headache Group - Department of Neurology, University of California, San Francisco,
San Francisco, CA, USA

³Department of Neuroscience, Amgen Inc.,
Thousand Oaks, CA, USA

Current address

[¶]Department of Neural and Pain Sciences, University of Maryland Baltimore, Baltimore, MD,
USA

*** These senior authors contributed equally to this manuscript**

Characters- Title: 87; Running Head: 26

Words- Abstract 186; Introduction 487; Discussion 141217; Total 59586050

Figures-7 (color- 2); Tables- 1

Number of pages: 46

Correspondence to:

Prof. Peter J. Goadsby
NIHR - Wellcome Trust Clinical Research Facility
King's College Hospital
London SE5 9PJ
UK
Email: peter.goadsby@kcl.ac.uk

Abstract

Pituitary adenylate cyclase activating polypeptide-38 (PACAP38) may play an important role in primary headaches. Preclinical evidence suggests that PACAP38 modulates trigeminal nociceptive activity mainly through PAC₁ receptors while clinical studies report that plasma concentrations of PACAP38 are elevated in spontaneous attacks of cluster headache and migraine and normalize after treatment with sumatriptan. Intravenous infusion of PACAP38 induces migraine-like attacks in migraineurs and cluster-like attacks in cluster headache patients.

A rodent-specific PAC₁ receptor antibody Ab181 was developed and its effect on nociceptive neuronal activity in the trigeminocervical complex was investigated *in vivo* in an electrophysiological model relevant to primary headaches.

Ab181 is potent and selective at the rat PAC₁ receptor and provides near maximum target coverage at 10 mg/kg for more than 48 hours. Without affecting spontaneous neuronal activity, Ab181 effectively inhibits stimulus-evoked activity in the trigeminocervical complex.

Immunohistochemical analysis revealed its binding in the trigeminal ganglion and sphenopalatine ganglion but not within the CNS suggesting a peripheral site of action.

The pharmacological approach using a specific PAC₁ receptor antibody could provide a novel mechanism with a potential clinical efficacy in the treatment of primary headaches.

Introduction

Migraine and cluster headache are highly disabling disorders [53] involving activation of the trigeminovascular system, which includes the perivascular meningeal afferents and the trigeminal ganglion in the peripheral nervous system, and the trigeminocervical complex (TCC) as well as higher brain structures including the hypothalamus, thalamus and periaqueductal gray in the central nervous system [18,34].

Pituitary adenylate cyclase activating polypeptide-38 (PACAP38) has received increasing attention in the context of migraine and cluster headache. PACAP38 is a 38-amino acid neuropeptide that is structurally and functionally related to vasoactive intestinal peptide (VIP) [27,40]. Both neuropeptides act on the same set of receptors, namely PAC₁, VPAC₁ and VPAC₂ receptors, which belong to the G-protein-coupled receptors of the secretin family [27]. PACAP38 [44] and VIP [22] have vasodilatory properties and play a major role in parasympathetic communication [7,27,36]. In cluster headache both neuropeptides are released during a spontaneous attack [15,52]. In migraine, despite some shared biology, both molecules have important differences. During a spontaneous migraine attack PACAP38 is released into the cranial circulation regardless of the presence of autonomic symptoms [51,55] while VIP is only released if cranial autonomic symptoms accompany the attack [17]. The infusion of PACAP38 [2,26,45], but less reliably VIP [43], induce migraine-like attacks in which can be effectively treated with sumatriptan. Sumatriptan normalizes elevated levels of PACAP38 during migraine [55]. These findings strikingly resemble the preclinical and clinical observations made with calcitonin gene-related peptide (CGRP) [14,16,17,37,57] which plays a prominent role in the pathophysiology of migraine and cluster headache and

has been proven to be a validated target in the treatment of both disorders
[13,18,24,30,39,47].

A large body of preclinical evidence has supported and begun to dissect the mechanisms behind these clinical observations. The fact that PACAP38 and VIP share similar affinities to the VPAC₁ and VPAC₂ receptors but that the PAC₁ receptor has a 100- to 1000-fold higher affinity to PACAP38 [9] has led to the conclusion that the relevant action of PACAP38 is likely to be mediated mainly through the PAC₁ receptor [55]. These findings are supported by *in vivo* studies that demonstrate the release of PACAP38 into the cranial circulation upon peripheral trigeminal activation [55,56] and the facilitation of nociceptive neuronal transmission in the TCC that is reversible by the administration of a PAC₁ receptor antagonist [1].

These observations suggest that targeting the PAC₁ receptor may offer an effective and highly selective approach to reduce trigeminal activation and offer a novel target for the treatment of primary headaches. We therefore developed a potent and selective monoclonal mouse anti-rat PAC₁ antibody (Ab181), fully characterized the pharmacologic and pharmacokinetic properties of the agent, and studied its effect on nociceptive neuronal activity within the TCC in an *in vivo* model that has been proven to be highly predictive for clinical efficacy in primary headaches [19-21,50]. Preliminary results have been presented at the 5th European Headache and Migraine Trust International Congress [32] and the International Headache Congress [33].

Methods

The generation and *in vitro* characterization of Ab181, the pharmacodynamic and pharmacokinetic analyses as well as the immunohistochemistry were conducted between 2009 and 2013 in the Amgen laboratories in Thousand Oaks, CA, USA. The electrophysiological studies were conducted between 2012 and 2013 in the laboratory of the Headache Group at the Department of Neurology, University of California, San Francisco, CA, USA.

The generation of the Ab181

Mice anti-rat PAC₁-specific monoclonal antibodies were generated using a conventional immunization method. Five 4-6-week old hybrid 129xC57BL/6 mice (Charles River Laboratories, Hollister, CA, USA) received three rounds of immunizations with soluble rat PAC₁, chemically conjugated to Padre Peptides (Pan DR Helper T-cell epitopes). Mice were immunized SQ/IP with up to 50 µg of antigen mixed or emulsified in either Freund's complete adjuvant (Cat# 77140, Pierce, Thermo Fisher Scientific, Waltham, MA, USA), RIBI (Cat# S6322, Sigma-Aldrich, St. Louis, MO, USA) or Poly I: C/CpG. Soluble PAC₁ receptor polypeptides containing the N-terminal extracellular domains (ECDs) of rat PAC₁ (amino acids 1-135 of GenBank accession no. NM133511.1) were generated by transiently cotransfecting 293-6E cells. All mice were maintained according to the regulations of the Amgen Institutional Animal Care and Use Committees (IACUC) in Thousand Oaks, CA. Mice with the highest detected FACS titer to Rat PAC₁ expressed on CHO AMID cells were selected for fusion. Four days prior to spleen harvest, mice selected for fusion were given a final IP boost of 50 µg soluble rat PAC₁ in PBS (Cat# 14040, GIBCO, Thermo Fisher Scientific, Waltham, MA, USA). B cell hybridomas were obtained by fusing immune splenocytes with

nonsecreting murine myeloma cells, Sp2/0-Ag14 (American Type Culture Collection), at a ratio of 2.5:1 by electrofusion. Ab181 is identified through screening assays including binding competition, functional blocking, and receptor selectivity against the rat PAC₁ receptor.

***In vitro* characterization of A**Ab**181**

Potency and selectivity of A**Ab**181 were analyzed *in vitro* cell-based functional assay.

Cell culture

In house developed stable rPAC₁/CHO cells were grown in Ham's F12 nutrient mixture (Cat# 11765, GIBCO, Thermo Fisher Scientific, Waltham, MA, USA); 10% fetal bovine serum (FBS) (Cat# 10099, GIBCO, Thermo Fisher Scientific, Waltham, MA, USA); 1X Penicillin-Streptomycin-Glutamine (Cat# 10378, GIBCO, Thermo Fisher Scientific, Waltham, MA, USA); 400 µg/mL G418 (Cat# 10131, GIBCO, Thermo Fisher Scientific, Waltham, MA, USA); 250 µg/ml Zeocin (Cat# R250-01, Invitrogen, Thermo Fisher Scientific Waltham, MA, USA). All cell flasks were maintained in incubators at 37°C with 5% CO₂. U2OS (Cat# HTB-96™, ATCC, Manassas, VA, USA) cells were grown in the medium of McCoy's 5A (Cat# 16600, GIBCO, Thermo Fisher Scientific, Waltham, MA, USA) containing 10% FBS (Cat# 10099, GIBCO, Thermo Fisher Scientific, Waltham, MA, USA), 1x L-glutamine (Cat# 25030, GIBCO, Thermo Fisher Scientific, Waltham, MA, USA), 1X Penicillin-Streptomycin-Glutamine and 1 X MEM Non-Essential Amino Acids (Cat# 11140, GIBCO, Thermo Fisher Scientific, Waltham, MA, USA). All cell flasks were maintained in incubator at 37°C with 5% CO₂.

BacMam virus construct

The rVPAC₁ and rVPAC₂ BacMam virus constructs were prepared in house. The titer of rVPAC₁ BacMam virus is 7.01×10^8 IU/ml, and the titer of rVPAC₂ BacMam virus is 5.75×10^8 IU/ml.

Preparation of BacMam virus transduced rVPAC₁ and rVPAC₂ cells:

U2OS cells were cultured in T-75 flask until the cell density reach to 70 - 80% confluent before transduction. Cell medium was removed from flask and the cells were rinsed with 1 X PBS (Cat# 14040, GIBCO, Thermo Fisher Scientific, Waltham, MA, USA) once, then Versene (Cat# 15040, GIBCO, Thermo Fisher Scientific, Waltham, MA, USA) was added to detach the cells. 3×10^6 U2OS cells were resuspended with culture medium and mixed with rVPAC₁ or rVPAC₂ BacMam virus at a concentration of 50 multiplicity of infection (MOI)/cell. The cell mixture was further incubated overnight for assay.

Cell-based functional assay

The cAMP assay was performed by using LANCE[®] cAMP ultra assay kit (Cat # TRF0263, PerkinElmer Inc., Waltham, MA, USA) to determine the activity of Ab181. Assay buffer contain Ham's F12 nutrient mixture (Cat# 11765, GIBCO, Thermo Fisher Scientific, Waltham, MA, USA), 0.1% bovine serum albumin (BSA) (Cat# CR84-100, PerkinElmer, Inc., Waltham, MA, USA), 1 mM IBMX (Cat# I5879, Sigma-Aldrich, St. Louis, MO, USA). Agonist dose response curve was first run to determine the appropriate concentration to be used in subsequent antagonism studies (data not shown) (Figure 1 and Table 1).

The antagonist activity of Ab181 was carried out by using EC₈₀ of PACAP38 (Cat# H-8430, Bachem, Bubendorf, Switzerland) or Maxadilan (Cat# H-6734, Bachem, Bubendorf,

Switzerland). PACAP6-38 (Cat# H-2734, Bachem, Bubendorf, Switzerland), a PAC₁ receptor antagonist, was used as a positive control. Ab181 (0.5 pM – 1 µM) was pre-incubated with the rPAC1-CHO cell suspension (2,000 cells/well) at room temperature for 30 min prior to the addition of agonists. PACAP38 or maxadilan was then added and incubated with the mixture for additional 15 min at room temperature. The reaction was stopped by adding detection mix of Eu-cAMP tracer and ULightTM-anti-cAMP (Cat# TRF0263, PerkinElmer Inc., Waltham, MA, USA) to all wells. After a 45-min incubation at room temperature, the assay plate was read on an EnVision instrument (Cat# 2105-0100, Perkin-Elmer Inc., Waltham, MA, USA).

Same assay method was used to determine the activity of Ab181 on rVPAC₁ and rVPAC₂ receptors. The activity of VIP (Cat#: H-3775, Bachem, Bubendorf, Switzerland), a rVPAC₁ and rVPAC₂ receptor agonist, was first evaluated and the EC₈₀ concentration of VIP was used in the assay to determine the activity of Ab181 on these receptors.

All results were analyzed using GraphPad Prism software's nonlinear regression curve fit (GraphPad Software, Inc., La Jolla, CA, USA) and data are presented as mean ± SD. Data from the agonist dose-response curves were used to calculate the half maximal effective concentration (EC₅₀) and the half maximal inhibitory concentration (IC₅₀) values for agonist and antagonist studies, respectively.

Pharmacokinetic and pharmacodynamic analysis

Male naïve Sprague Dawley rats, 6-12 weeks from either Taconic Farms Inc. (Oxnard, CA, USA) or Harlan Laboratories (Indianapolis, IN, USA) at the average age of initiation of treatment were used and all procedures in this report were conducted in compliance with

the Animal Welfare Act, the Guide for the Care and Use of Laboratory Animals, and the Office of Laboratory Animal Welfare. Animals were group-housed in non-sterile, ventilated micro-isolator housing on corn cob bedding in Amgen's Assessment and Accreditation of Laboratory Animal Committee (AAALAC)-accredited facility with controlled temperature ($70 \pm 5^\circ \text{F}$), relative humidity ($50 \pm 20\%$), and 12-h light/dark cycles (0600 to 1800). Animals had *ad libitum* access to pelleted feed (Harlan Teklad 2020X, Indianapolis, IN, USA) and water (on-site generated reverse osmosis) via automatic watering system.

Test and control materials

Ab181 was generated from Amgen and diluted to a series of concentrations in A5Su (10 mM sodium acetate, 9% sucrose, pH = 5.0). An isotype control was used as a dummy antibody in A5Su at a concentration of 5 mg/ml. Maxadilan (trifluoroacetate salt, Bachem, Bubendorf, Switzerland) was used and a dosing solution was freshly prepared daily by dissolving maxadilan in 1X DPBS (Dulbecco's phosphate buffered saline) (Sigma-Aldrich, St. Louis, MO, USA) at a final concentration of 0.5 $\mu\text{g/ml}$.

Laser Doppler Imaging

A laser Doppler imager (Moor Instruments, Ltd, Wilmington, DE, USA) was used to measure DBF on the shaved skin of the rat abdomen.

Following anesthetic with propofol on the test day, the rat's abdominal area was shaved and each animal was placed in a supine position on a temperature-controlled circulating warm-water heating pad to help maintain a stable body temperature during the study. After a 10 to 15-min stabilization period, a black rubber O-ring (0.925 cm inner diameter, O-Rings West, Seattle, WA, USA) was placed on the rat abdomen without directly positioning it over

1 a visible blood vessel. After placement of an O-ring on the selected area, a baseline (BL) DBF
2 measurement was taken. After the BL scan, the maxadilan solution prepared fresh daily in
3
4 20 µl vehicle (DPBS) was injected intradermally at the center of the O-ring. The post-
5
6
7 maxadilan DBF was measured either every 15 min over a 60-min period or at specified time
8
9
10 points such as 15 and 30 min. The O-ring serves as an area of interest in which the DBF will
11
12
13 be analyzed within the O-ring. Ab181 was prepared in A5Su at different concentrations
14
15 depend on the dose range and given in a single bolus i.v. injection. In this report, MIIBF was
16
17
18 measured and expressed as % change from the baseline [$100 \times (\text{individual post-agonist flux} -$
19
20
21 individual baseline flux)/individual baseline flux] or as % inhibition [(Mean of % change from
22
23
24 BL from vehicle-treated animals - individual % change from BL from drug treated animals)/
25
26
27 Mean % change from BL from vehicle-treated animals].
28
29
30

31 *Dose-response study at 48 h post Ab181 treatment*

32
33 Ab181 was administered via penile vein or tail vein at various doses (0.1, 0.3, 1 and 10
34
35 mg/kg). Forty-eight h later, rats received an intradermal maxadilan injection (10 ng in 20 µl
36
37
38 DPBS) followed by post-maxadilan DBF scans every 15 min over a 60-min period. Following
39
40
41 the final DBF scan, three rats that had been pre-treated with 10 mg/kg of Ab181 were
42
43
44 allowed to recover from the propofol anesthesia, returned to their home cages and
45
46
47 underwent a second DBF measurement at 168 h post-drug administration. Serum PK
48
49
50 samples were taken via tail vein immediately prior to the first maxadilan challenge at 48 and
51
52
53 168 h post Ab181 treatment. The post MIIBF scan time point for the later studies was
54
55
56 determined based on the post-maxadilan DBF response from 15 to 60 min.
57
58

59 *Dose-response study at 3.25 h post Ab181 treatment*

60
61
62
63
64
65

Ab181 was administered via the penile vein at various doses (0.1, 0.3, 1, 3 and 10 mg/kg).

Three hours later, rats received an intradermal maxadilan injection (10 ng of maxadilan in 20 μ l DPBS) followed by a post-MIBF scan 15 min later. Serum PK samples were taken via tail vein immediately after the DBF scan (i.e., 3.25 h post Ab181 treatment). [The time point was chosen to confirm that the receptor occupancy of Ab181 had reached and stayed at the desired pharmacodynamic measures when performing the electrophysiological studies.](#)

Time-course study of Ab181 at a dose of 10 mg/kg

Ab181 was dosed intravenously at 10 mg/kg via either rat tail vein or penile vein at various pre-treatment times (0.5, 3, 6.25 h) prior to maxadilan challenge. The full time-course study was comprised of several experiments, including the 48- and 168 h post-Ab181 treatment studies described above.

Statistical analysis of laser Doppler flow experiments

All DBF results were expressed as the mean \pm SEM. A one-way ANOVA followed by Dunnett's Multiple Comparison Test (MCT) was used to assess the statistical significance of Ab181 effects relative to either vehicle or the control antibody. A $p < 0.05$ was used to determine significance between any two groups. ED₅₀ values were calculated in GraphPad Prism following logarithmic transformation and nonlinear fit of data using a sigmoidal dose-response model with variable slope and with the top constrained to the resulting mean percent change in the corresponding control group (vehicle or control antibody-treated group) and bottom constrained to zero.

Electrophysiological recordings

All experiments were approved by the Institutional Animal Care and Use Committee (IACUC) of the University of California, San Francisco. Experiments were conducted in accordance with the United States Public Health Service's Policy on Humane Care and Use of Laboratory Animals, the ARRIVE guidelines and the guidelines of the Committee for Research and Ethical Issues of the International Association of the Study of Pain (IASP).

General surgical preparation

Twenty-four male Sprague-Dawley rats (Charles River Laboratories, Hollister, CA, USA) were used in the experiments. [In each animal only one experiment was conducted and recordings were performed from one site.](#) The animals were anesthetized by an induction with a single dose of pentobarbital (60 mg/kg i.p.; Nembutal, Lundbeck, Deerfield, IL, USA) followed by a continuous infusion of propofol (20-25 mg/kg/h i.v.; Propoflo, Abbott, Abbott Park, IL, USA) for maintenance throughout the entire experiment. For the administration of the anesthetic and drugs both femoral veins were cannulated. The left femoral artery was cannulated for the continuous monitoring of arterial blood pressure.

Physiological monitoring

Rats were placed on a self-regulating homeothermic blanket system with a rectal probe (Harvard Apparatus, Holliston, MA, USA) and core body temperature was maintained at 37 ± 0.5 °C. Arterial blood pressure was monitored from the femoral artery using a transducer (DTX Plus DT-XX, Becton Dickinson, Sandy, UT, USA) connected to an amplifier (PM-1000, CWE, Ardmore, PA, USA). Following a tracheostomy, animals were mechanically ventilated (3-5 ml/min, 75-90 strokes/min; 7025, Ugo Basile, Comerio, VA, Italy) with oxygen-enriched air and end-expiratory CO₂ was kept between 3.5 and 4.5%. Data on arterial blood pressure

and CO₂ concentration were continuously displayed and fed into a data acquisition system (Power 1401, Cambridge Electronic Design-CED, Cambridge, UK) and saved on a hard disk.

Recording preparation

The rat's heads were fixed in a stereotaxic frame (Kopf Instruments, Tujunga, CA, USA). A craniotomy was performed over the parietal cortex with a dental burr using constant irrigation to reduce heat production. With this procedure the middle meningeal artery (MMA) was exposed without lesioning the dura mater. For electrical stimulation a bipolar stimulating electrode (NE200, Rhodes Medical Instruments, Summerland, CA, USA) was placed above the MMA touching the dura mater at either side of the blood vessel. The electrode was connected to a stimulus isolation unit (SIU5A, Grass Instruments, Quincy, MA, USA).

For the extracellular recording of neuronal activity in the trigeminocervical complex (TCC) a C1 partial hemilaminectomy was performed and the spinal dura mater was removed. A tungsten electrode with a nominal impedance of 1 M Ω (TM31A10, World Precision Instruments, Sarasota, FL, USA) was then introduced in the TCC near the dorsal root entry zone. For the localization of optimal site for the extracellular recording, the electrode was advanced or retracted in 5 μ m steps with a piezoelectric motor-driven micromanipulator. Wide dynamic range (WDR) neurons with convergent input from the dura mater and the facial skin were identified by their responsiveness to electrical stimulation of the perivascular afferents surrounding the MMA as well as innocuous brush and noxious pinch of the skin innervated by the first branch of the trigeminal nerve.

Stimulation of meningeal afferents and recording in the TCC

Electrical stimulation of the perivascular meningeal afferents was performed applying electrical square wave pulses (10-18 V, 0.1-0.2 ms, 0.5 Hz, 20 sweeps) (S88, Grass Instruments, Quincy, MA, USA).

The stimulus-evoked neuronal signal as well as the neuronal background activity were acquired by the recording electrode placed in the TCC. The electrical signal was fed into a headstage amplifier (NL100AK, Neurolog, Digitimer Welwyn Garden City, Hertfordshire, UK) and passed to an AC preamplifier (NL104, Neurolog, Digitimer Welwyn Garden City, Hertfordshire, UK) set to a gain of 1000x. The signal was then passed through a band-pass filter (bandwidth 300 Hz to 10 kHz) (NL125/126, Neurolog, Digitimer Welwyn Garden City, Hertfordshire, UK) and a 60 Hz noise eliminator (Humbug, Quest Scientific, Vancouver, BC, Canada) before further amplification by an AC-DC amplifier (NL106, Neurolog, Digitimer Welwyn Garden City, Hertfordshire, UK). This signal was fed to a gated amplitude discriminator (NL201 Neurolog, Digitimer Welwyn Garden City, Hertfordshire, UK) and a data acquisition system (Power 1401, Cambridge Electronic Design-CED, Cambridge, UK). Data was collected, analyzed and stored using Spike 5.2 software (Cambridge Electronic Design-CED, Hertfordshire, UK). The output of the gated amplitude discriminator was also fed into an audio amplifier (NL120, Neurolog, Digitimer Welwyn Garden City, Hertfordshire, UK) and loudspeaker as well as an oscilloscope to assist spike discrimination from background activity. For the analysis of stimulus-evoked neuronal activity post-stimulus histograms were produced online. Background activity gated through the amplitude discriminator was collected into successive bins.

Experimental protocol and drug administration

After completing the surgical procedure, the animals had a resting period of 30 min.

Following this period, baseline recordings were obtained. These were obtained by calculating the mean of 3 series of post-stimulus histograms, each consisting of 20 electrical stimuli.

Following the assessment of the baseline values Ab181 (10 mg/kg) or its vehicle (A5Su) were administered intravenously over 1 min. Due to the pharmacological properties of the antibody the animals underwent a second resting period of 2.5 h to allow sufficient time for the antibody to bind its target. After this second resting sumatriptan (10 mg/kg) or vehicle were administered intravenously over 1 min followed by another resting period of 30 min. Post-stimulus histograms were then established 180, 185, 190, 195, 200, 205, 210 and 225 min after the administration of the first pharmacological intervention (Ab181 or vehicle) (Figure 3.1). Based on the treatments described above, animals were divided in 3 treatment groups, group 1 receiving the Ab181 (intervention 1) and sumatriptan vehicle (intervention 2), group 2 receiving vehicle (intervention 1) and sumatriptan (intervention 2) and group 3 receiving vehicle at both interventions (Figure 3.1).

Statistical analysis

Statistical analysis was performed using SPSS 22.0 software (IBM Corporation, Armonk, NY, USA). Data are expressed as percentages of baseline values with standard errors of the mean (\pm SEM). Effects within a treatment group were calculated using the analysis of variance (ANOVA) with repeated measures applying the Greenhouse-Geisser correction of the assumption of sphericity was violated. Bonferroni correction was applied for multiple comparisons. Statistical significance was assumed at $p < 0.05$. For a detailed comparison of

individual data points with the baseline value within one treatment group the dependent *t*-test was used.

Immunohistochemistry

Adult male Sprague Dawley rats (n = 3 per group) were injected intravenously with Ab181 or a monoclonal mouse isotype control antibody against an unrelated target or vehicle. After 3.5 h, animals were terminally anesthetized by FatalPlus™ (Vortech Pharmaceuticals, Dearborn, MI, USA) and perfused with ice cold phosphate-buffered saline (PBS) at pH 7.4 (Cat# 14040, GIBCO, Thermo Fisher Scientific, Waltham, MA, USA) followed by 4% paraformaldehyde (Cat# P6148, Sigma-Aldrich, St. Louis, MO, USA) in PBS (Cat# 14040, GIBCO, Thermo Fisher Scientific, Waltham, MA, USA). The SPG, TG and brains were dissected and cryoprotected with 30% sucrose in PBS. 12 µm sections were cut, washed in PBS and incubated in 3% normal goat serum (NGS) and 0.3% TritonX-100 in PBS for one hour at room temperature. Sections were then washed in PBS and incubated with AlexaFluor™488 goat anti-mouse IgG (Invitrogen, Thermo Fisher Scientific Waltham, MA, USA) for 1 h at room temperature. After final washes in PBS (Cat# 14040, GIBCO, Thermo Fisher Scientific, Waltham, MA, USA), sections were cover slipped with Vectashield mounting medium with DAPI (Cat# H-1200, Vector Labs, Burlingame, CA, USA). The following additional controls were conducted: A set of sections from Ab181-dosed rats was incubated with secondary antibodies against rabbit (AlexaFluor™488 goat anti-rabbit IgG, Invitrogen, Thermo Fisher Scientific, Waltham, MA, USA) as negative control. As positive control for PAC₁ staining a set of vehicle sections was incubated with 10 µg/ml Ab181 in blocking solution at 4 °C for 48 h before further processing with AlexaFluor™488 goat anti-mouse IgG (Invitrogen, Thermo Fisher Scientific Waltham, MA, USA) as described above.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Results

In vitro potency of the PAC₁ receptor antibody Ab181

Ab181 is a full antagonist of the rat PAC₁ receptor. It dose-dependently inhibited PAC₁ receptor agonist PACAP38 or maxadilan, a PAC₁ selective receptor agonist, induced cAMP accumulation in CHO cells expressing rat PAC₁ receptors with IC₅₀ of 20 ± 3.3 nM ($n = 3$) and 4.5 ± 0.1 nM ($n = 2$), respectively (Figure 1, Table 1). Ab181 is selective to the PAC₁ receptor. IC₅₀ at rat VPAC₁ and rat VPAC₂ receptors against the agonist VIP is greater than 100 nM, the highest concentration tested.

Pharmacokinetic and pharmacodynamic effect of Ab181

Dose-response study at 48 h post Ab181 treatment

Pretreatment of Ab181 48 h prior to maxadilan challenge prevented the maxadilan-induced increase in dermal blood flow (MIIDBF). The change in dermal blood flow (DBF) over 60 min post-maxadilan is depicted in Figure 2.1A. There was a statistically significant difference ($p < 0.001$) between the vehicle-treated group and each of three Ab181 dose groups (0.3, 1 and 10 mg/kg) at 15 min post maxadilan injection with a calculated ED₅₀ of 0.16 mg/kg (CI_{95%} = 0.10-0.26 mg/kg). Serum concentration of Ab181 is plotted on the Y-axis of Figure 2.1B.

Dose-response study at 3.25 h post Ab181 treatment

A shorter pretreatment of Ab181 at various doses was also evaluated. A fifteen minute post-maxadilan measurement was used in this study. The change in MIIDBF after 3.25 h post-Ab181 is depicted in Figure 2.2. There was a statistically significant inhibition of the MIIDBF following dose of 1, 3 and 10 mg/kg at 3.25 h post treatment ($p < 0.01$), resulting in a

calculated ED₅₀ of 0.95 mg/kg (CI_{95%} = 0.59-1.51 mg/kg). Serum concentration of Ab181 is plotted on the Y-axis of Figure 2.2.

Time-course study of Ab181 at a dose of 10 mg/kg

A fifteen minute post-maxadilan measurement was used in this study. Ab181 at 10 mg/kg produced a statistically significant inhibition of the MIIDBF starting ($p < 0.05$) from a pretreatment time of 0.75 h and lasting through the last measured 168 h time point (Figure 2.3). The resulting mean serum concentration at 1, 3, 6, 48 and 168 h post drug treatment is plotted on the Y-axis in Figure 2.3.

Electrophysiological recordings

Ab181 inhibits stimulus-evoked responses in the TCC

The intravenous administration of Ab181 (Figure 3.1) induced a long-lasting inhibition of stimulus-evoked nociceptive neuronal activity within the TCC ($F_{1.48, 5.91} = 8.43$, $p = 0.022$). Compared to the baseline value, the effect was significant at all investigated time points throughout the entire observational period. Maximum inhibition reached $-43 \pm 13\%$ ($t_7 = 3.41$, $p = 0.011$) compared to baseline.

Sumatriptan, which was administered as a positive control, significantly reduced stimulus-evoked responses in the TCC ($F_{2.21, 15.49} = 4.97$, $p = 0.019$). Similar to the Ab181 group, the inhibiting effect of sumatriptan was significant at all investigated time points reaching a maximum inhibition of $-42 \pm 15\%$ ($t_7 = 2.78$, $p = 0.027$) compared to baseline without recovering until the end of the observational period.

In contrast, the intravenous administration of the vehicle did not affect stimulus-evoked neuronal activity ($F_{2.21, 15.48} = 2.45$, $p = 0.115$; Figure 3.2A, B).

Neuronal background activity in the TCC is not affected by Ab181

In contrast to the observed effect on stimulus-evoked neuronal activity, intravenous administration of Ab181 does not attenuate unspecific neuronal background activity within the TCC ($F_{1.32, 5.27} = 0.767$, $p = 0.456$). The same lack of effect on background activity was observed in animals treated with sumatriptan ($F_{2.33, 16.28} = 0.821$, $p = 0.474$) and those treated with vehicle ($F_{1.68, 11.78} = 0.99$, $p = 0.385$; Figure 3.2C).

Arterial blood pressure is unaffected by Ab181

Intravenous administration of Ab181 does not affect arterial blood pressure ($F_{1.69, 6.78} = 0.656$, $p = 0.525$). In contrast, sumatriptan induced a significant decrease in arterial blood pressure ($F_{2.55, 17.84} = 5.82$, $p = 0.008$) throughout the entire observational period, reaching a maximum reduction of $-13 \pm 3\%$ ($t_7 = 4.89$, $p = 0.002$) compared to baseline without recovering until the end of the experiment. In the vehicle control group arterial blood pressure was unaffected ($F_{2.29, 16.02} = 0.93$, $p = 0.426$; Figure 3.2D).

Immunohistochemistry

To investigate whether Ab181 or the control antibody distributed into tissues of interest after intravenous administration, we treated satellite rats from the electrophysiology study, but dissected and fixed tissues of interest at the 3.5 h time point after injection. Using fluorescently labeled secondary antibodies against the Fc-portion of the antibodies we were able to detect immunoreactivity in dura, trigeminal (TG) and sphenopalatine ganglia (SPG) of

Ab181-dosed rats, but not in rats dosed with control antibody (Figure 4). In contrast, no labeling above background was detected in spinal trigeminal nucleus or superior salivary nucleus of the brainstem as well as in the hypothalamus and thalamus indicating that Ab181 did not cross the blood-brain-barrier or the amount of antibodies that entered the CNS after intravenous injection was below the detection threshold of the immunohistochemical method (Figure 4).

Discussion

PACAP38, first described in 1989 [40], is one of the key neurotransmitters of the parasympathetic system. Soon after its discovery, preclinical evidence suggested effects beyond its parasympathetic role, in particular in specific pain pathways related to headache [56]. Neuroanatomical and functional data obtained since strongly suggests a prominent role in primary headaches including cluster headache and migraine. Our data [may](#) provide a plausible under-pinning biology for a [potential](#) clinical effect of PAC₁ blockade in these disorders.

First, PACAP receptors (PAC₁, VPAC₁ and VPAC₂) are located in several key areas along the trigeminovascular system. Within the central part of the pain pathways processing trigeminal pain they have been identified in the thalamus [28], hypothalamus [28] and the TCC [35] while in the peripheral part, they are located in the TG [35] and the meningeal vasculature [5,6]. Beyond that, they are found in the SPG [8] highlighting the relevance of PACAP38 in the parasympathetic system, as well as its potential role in the functional interaction between the trigeminal and parasympathetic systems, namely the trigeminoautonomic reflex.

Secondly, from a functional perspective, PACAP38 may play a role in cluster headache and migraine since its plasma concentration is elevated in spontaneous attacks [52,55] and normalizes after effective attack-abortion with sumatriptan [55]. In addition, PACAP38 triggers migraine-like attacks in migraineurs [26,44], most likely through an activation of the PAC₁ receptor [55]. In the context of migraine, the ability to trigger attacks clearly distinguishes PACAP38 from VIP. These clinical observations can now be explored in specific

1 animal models. For example, electrical stimulation of the superior sagittal sinus of the cat
2 induces the release of PACAP38 into the cranial circulation [55,56]. In line, the
3
4 administration of PACAP38 facilitates nociceptive neuronal activity in the TCC [1]. These
5
6 functional features of PACAP38 strikingly resemble the preclinical [50] and clinical
7
8 observations [16,17] made with CGRP which predicted clinical efficacy of six small molecule
9
10 CGRP receptor antagonists [10,29,31,38,42,54] and CGRP/CGRP-receptor antibodies
11
12 [11,23,46,48] in the acute and preventive treatment of migraine as well as a CGRP antibody
13
14 in the preventive treatment of episodic cluster headache [13].
15
16
17
18
19
20
21
22

23 Thirdly, the parasympathetic and trigeminal sensory systems connect and interact with each
24
25 other. From an anatomic perspective, CGRP-positive sensory fibers project from the TG to
26
27 the SPG [8] and *in vivo* data demonstrates that PACAP38 induces the release of CGRP from
28
29 the TCC [35]. Nevertheless, the detailed molecular mechanisms behind the attack-triggering
30
31 capability of PACAP38 and in particular the clinical relevance of the above described
32
33 mechanism remain to be fully elucidated. Interestingly, the administration of PACAP38,
34
35 despite creating an activation of the trigeminovascular system with the clinical picture of a
36
37 migraine-like attack, does not induce the release of CGRP in a human model of migraine
38
39 [26]. These results suggest that the PACAP38-induced effects on trigeminal activation are
40
41 likely to be the result of a direct effect on nociceptive trigeminal neurons rather than a
42
43 CGRP-mediated mechanism although both mechanisms do not exclude each other. In the
44
45 case of a direct activation of either of the PACAP receptors, PACAP38 stimulates the
46
47 cAMP/PKA pathway thereby increasing neuronal excitability [49]. In addition, all PACAP
48
49 receptors may be modulated in their expression profile under chronic pain conditions [58].
50
51 It may be speculated that these up- and down regulating properties and the ability to
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 produce neuronal sensitization may even play a role in the chronification of migraine. Based
2 on this large body of evidence, we set out to develop a specific antibody against the PAC₁
3 receptor as this receptor, as outlined above, is most likely the most relevant PACAP receptor
4 in mediating the attack-triggering effect of PACAP38 and to test its effects on a model
5 system that represents significant elements of cluster headache and migraine
6 pathophysiology [4].
7
8
9
10
11
12
13
14
15
16
17

18 PACAP38 is a non-selective agonist at all the PACAP receptors, and VIP is only mildly
19 selective to the VPAC₁ and VPAC₂ receptors [27]. Maxadilan, on the other hand, is the most
20 selective agonist at the PAC₁ receptor [41], which allowed us to measure pharmacodynamic
21 activities specifically mediated through the PAC₁ receptor. Although some years have passed
22 since PACAP38 was discovered [40], selective antagonists to the individual receptors in the
23 PACAP receptor family are still lacking. Several peptide antagonists have been reported with
24 mild selectivity between different receptors [5,9,25] but their pharmacokinetic properties,
25 especially their very short plasma half-life, hindered their utility in *in vivo* pharmacology
26 studies. Therefore, Ab181 was developed. As demonstrated in the Results section, it is a
27 potent and selective antagonist at the PAC₁ receptor, with no activity at the VPAC receptors.
28 It also potent and selective inhibitor of maxadilan-induced increases in dermal blood flow in
29 a time- and dose-dependent manner. Through the *in vitro* and *in vivo* profiling, Ab181
30 demonstrated full target coverage of the PAC₁ receptor at 3.5 h after intravenous
31 administration and maintained sustained plasma concentration throughout the study
32 duration. It is therefore the first ideal tool for the studying of selective PAC₁ pharmacology
33 in preclinical species.
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 The new results of the study show that the intravenous administration of Ab181 inhibits
2 stimulus-evoked nociceptive activity in the TCC. Unlike conventional small molecular agents
3 whose molecule weight (MW) are normally < 500, the much larger sized antibodies (MW >
4 145KD) generally have a slower tissue distribution rate. As a result, it requires a substantial
5 equilibrium period between the administration of the antibody and the initiation of the
6 experiment before the assessment of its influence on stimulus-evoked activity. Therefore,
7 the experimental design resembles more of a short-term preventive scenario than an acute
8 reversal of nociceptive trigeminal activation. The results show that the extent of neuronal
9 inhibition was almost identical to that observed with sumatriptan, suggesting a robust
10 effect. The fact that we did not observe an effect on non-specific background activity
11 reflects the specificity of the effect on nociceptive neuronal transmission.

~~12 and, beyond that, may also suggest a purely (or mainly) peripheral site of action.~~

13 The exact site of action of PAC₁ remains speculative. Literature suggested that agonist
14 PACAP38, with MW >4000, triggers migraine-like attacks without crossing the blood-brain-
15 barrier [12]. However, passage of PACAP27 and PACAP38 across the blood-brain barrier has
16 been suggested [3]. Here, immunohistochemical analysis of Ab181 found a minimum
17 presence in CNS which suggests that the site of the modulating action on nociceptive
18 activity is likely to be located outside of the CNS. This would be in line with the results
19 obtained from the electrophysiological experiments showing no effect on neuronal
20 background activity as well as the immunohistochemical analysis in which Ab181 was
21 detected in the relevant peripheral structures but not within the CNS. These findings
22 suggest that Ab181 does not cross the blood-brain barrier in a significant concentration, a
23 finding that is expected given the molecular size of the antibody. It cannot be excluded that

1 a small amount of antibody reaches the CNS but remains below the immunohistochemical
2 detection limit, however, target coverage with such small amount is unlikely sufficient for a
3 full antagonism effect. Therefore, the current study with a selective PAC₁ antagonist
4 antibody supports the hypothesis that peripheral PAC₁ receptor inhibition can be sufficient
5 to abort or prevent attacks of cluster headache and migraine.
6
7
8
9
10
11
12
13
14

15 Recently, a randomized, placebo-controlled trial (Phase 2) with a monoclonal PAC₁ receptor
16 antibody (AMG 301) was reported in abstract form to be ineffective in the preventive
17 treatment of migraine. In the context of our data some comments are relevant. First, in our
18 experiments we used an antibody that was developed for rodents and differs significantly in
19 its pharmacological properties, including its affinity to the PAC₁ receptor, [for which it is](#)
20 [more potent than the one used in the human Phase 2 trial.](#) ~~from the antibody used in the~~
21 ~~human Phase 2 trial.~~ Secondly, while the available evidence suggests that the PAC₁ receptor
22 plays a relevant role in trigeminal activation, the possibility that VPAC receptors are
23 clinically significant is an open issue. The fact that VIP may trigger migraine attacks in subset
24 of migraineurs may support this hypothesis [2]. Finally, in the clinical trial patients were not
25 stratified by the presence or absence of cranial autonomic symptoms. Therefore, it may be
26 hypothesized that the ability of PACAP38 to induce migraine attacks may require an action
27 on PAC₁ receptors on trigeminal and parasympathetic neurons thereby increasing the
28 activation of the trigeminoautonomic reflex. If this would be the case one could speculate of
29 a higher relevance of this potentially therapeutic mechanism in migraine with cranial
30 autonomic symptoms or in cluster headache.
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 Taken together, the findings of our studies suggest that the PAC₁ antibody Ab181 can inhibit
2 nociceptive neuronal traffic so that this pharmacological approach may offer a new strategy
3 for the preventive treatment of primary headache disorders, including cluster headache and
4
5 migraine.
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Acknowledgments

The study was funded by Amgen Inc., Thousand Oaks, CA, USA.

Conflicts of interest

Jan Hoffmann is consulting for and/or serves on advisory boards of Allergan, Autonomic Technologies Inc. (ATI), Chordate Medical AB, Eli Lilly, Hormosan Pharma, Novartis and Teva. He has received honoraria for speaking from Allergan, Chordate Medical AB, Novartis and Teva. He received personal fees for MedicoLegal Work and from Quintessence Publishing.

Silke Miller is an employee of Amgen Inc.

Margarida Martins-Oliveira does not report any conflict of interest.

Simon Akerman reports an unrestricted grant, honoraria and travel reimbursements from electroCore LLC, unrelated to the submitted work.

Weera Suprongsinchai does not report any conflict of interest.

Hong Sun is an employee of Amgen Inc.

Licheng Shi is an employee of Amgen Inc.

Dawn Zhu is an employee of Amgen Inc.

Sonya Lehto is an employee of Amgen Inc.

Hantao Liu is an employee of Amgen Inc.

Ruoyuan Yin is an employee of Amgen Inc.

Bryan D. Moyer is an employee of Amgen Inc.

Cen Xu is an employee of Amgen Inc.

Peter Goadsby reports grants and personal fees from Amgen and Eli-Lilly and Company, and personal fees from Alder Biopharmaceuticals, Allergan, Autonomic Technologies Inc., Biohaven Pharmaceuticals Inc., Electrocore LLC, eNeura, Impel Neuropharma, MundiPharma, Novartis, Teva Pharmaceuticals, and Trigemina Inc. WL Gore, and personal fees from MedicoLegal work, Massachusetts Medical Society, Up-to-Date, Oxford University Press, and Wolters Kluwer; and a patent Magnetic stimulation for headache assigned to eNeura without fee.

References

- [1] Akerman S, Goadsby PJ. Neuronal PAC1 receptors mediate delayed activation and sensitization of trigeminocervical neurons: relevance to migraine. *Science Translational Medicine* 2015;7:1-11.
- [2] Amin FM, Hougaard A, Schytz HW, Asghar MS, Lundholm E, Parvaiz AI, de Koning PJH, Andersen MR, Larsson HBW, Fahrenkrug J, Olesen J, Ashina M. Investigation of the pathophysiological mechanisms of migraine attacks induced by pituitary adenylate cyclase-activating polypeptide-38. *Brain* 2014;137(3):779-794.
- [3] Banks WA, Kastin AJ, Komaki G, Arimura A. Pituitary adenylate cyclase activating polypeptide (PACAP) can cross the vascular component of the blood-testis barrier in the mouse. *J Androl* 1993;14(3):170-173.
- [4] Bergerot A, Holland PR, Akerman S, Bartsch T, Ahn AH, MaassenVanDenBrink A, Reuter U, Tassorelli C, Schoenen J, Mitsikostas DD, van den Maagdenberg AMJM, Goadsby PJ. Animal models of migraine. Looking at the component parts of a complex disorder. *European Journal of Neuroscience* 2006;24:1517-1534.
- [5] Boni LJ, Ploug KB, Olesen J, Jansen-Olesen I, Gupta S. The in vivo effect of VIP, PACAP-38 and PACAP-27 and mRNA expression of their receptors in rat middle meningeal artery. *Cephalalgia* 2009;29:837-847.
- [6] Chan KY, Baun M, de Vries R, van den Boogaert AJ, Dirven CM, Danser AH, Jansen Olesen I, Olesen J, Villalon CM, Maassen Van Den Brink A, Gupta S. Pharmacological

characterization of VIP and PACAP receptors in the human meningeal and coronary artery. *Cephalalgia* 2011;31:181-189.

[7] Couvineau A, Laburthe M. VPAC receptors: structure, molecular pharmacology and interaction with accessory proteins. *British Journal of Pharmacology* 2012;166:42-50.

[8] Csati A, Tajti J, Kuris A, Tuka B, Edvinsson L, Warfvinge K. Distribution of vasoactive intestinal polypeptide, pituitary adenylate cyclase activating peptide, nitric oxide synthase and their receptors in human and rat sphenopalatine ganglion. *Neuroscience* 2012;202:158-168.

[9] Dickson L, Finlayson K. VPAC and PAC receptors: From ligands to function. *Pharmacol Ther* 2009;121(3):294-316.

[10] Diener H-C, Barbanti P, Dahlof C, Reuter U, Habeck J, Podhorna J. BI 44370 TA, an oral CGRP antagonist for the acute treatment of migraine attacks: results from a phase II study. *Cephalalgia* 2011;31:573-584.

[11] Dodick DW, Goadsby PJ, Silberstein SD, Lipton RB, Olesen J, Ashina M, Wilks K, Kudrow D, Kroll R, Kohrman B, Bargar R, Hirman J, Smith J. Randomized, Double-blind, Placebo-controlled, Phase II Trial of ALD403, an anti-CGRP peptide antibody in the prevention of frequent episodic migraine. *Lancet Neurology* 2014;13:1100-1107.

- [12] Erdling A, Sheykhzade M, Maddahi A, Bari F, Edvinsson L. VIP/PACAP receptors in cerebral arteries of rat: characterization, localization and relation to intracellular calcium. *Neuropeptides* 2013;47(2):85-92.
- [13] Goadsby PJ, Dodick DW, Leone M, Bardos JN, Oakes TM, Millen BA, Zhou C, Dowsett SA, Aurora SK, Ahn AH, Yang J-Y, Conley RR, Martinez JM. Trial of Galcanezumab in Prevention of Episodic Cluster Headache. *New England Journal of Medicine* 2019;381(2):132-141.
- [14] Goadsby PJ, Edvinsson L. The trigeminovascular system and migraine: studies characterizing cerebrovascular and neuropeptide changes seen in humans and cats. *Annals of Neurology* 1993;33:48-56.
- [15] Goadsby PJ, Edvinsson L. Human *in vivo* evidence for trigeminovascular activation in cluster headache. *Brain* 1994;117:427-434.
- [16] Goadsby PJ, Edvinsson L, Ekman R. Release of vasoactive peptides in the extracerebral circulation of man and the cat during activation of the trigeminovascular system. *Annals of Neurology* 1988;23:193-196.
- [17] Goadsby PJ, Edvinsson L, Ekman R. Vasoactive peptide release in the extracerebral circulation of humans during migraine headache. *Annals of Neurology* 1990;28:183-187.

- [18] Goadsby PJ, Holland PR, Martins-Oliveira M, Hoffmann J, Schankin C, Akerman S.
Pathophysiology of Migraine- A disorder of sensory processing. *Physiological Reviews*
2017;97:553-622.
- [19] Goadsby PJ, Hoskin KL. Inhibition of trigeminal neurons by intravenous administration
of the serotonin (5HT)_{1B/D} receptor agonist zolmitriptan (311C90): are brain stem
sites a therapeutic target in migraine? *Pain* 1996;67:355-359.
- [20] Goadsby PJ, Hoskin KL. Serotonin inhibits trigeminal nucleus activity evoked by
craniovascular stimulation through a 5-HT_{1B/1D} receptor: a central action in migraine?
Annals of Neurology 1998;43:711-718.
- [21] Goadsby PJ, Hoskin KL, Knight YE. Substance P blockade with the potent and centrally
acting antagonist GR205171 does not effect central trigeminal activity with superior
sagittal sinus stimulation. *Neuroscience* 1998;86:337-343.
- [22] Goadsby PJ, Macdonald GJ. Extracranial vasodilatation mediated by VIP (Vasoactive
Intestinal Polypeptide). *Brain Research* 1985;329:285-288.
- [23] Goadsby PJ, Reuter U, Hallstrom Y, Broessner G, Bonner JH, Zhang F, Sapra S, Picard H,
Mikol DD, Lenz RA. A controlled trial of erenumab for episodic migraine. *New*
England Journal of Medicine 2017;377:2123-2132.

- [24] Goadsby PJ, Reuter U, Hallstrom Y, Broessner G, Bonner JH, Zhang F, Sapra S, Picard H, Mikol DD, Lenz RA. A Controlled Trial of Erenumab for Episodic Migraine. *N Engl J Med* 2017;377(22):2123-2132.
- [25] Gourlet P, Vandermeers A, Vandermeers-Piret MC, Rathe J, De Neef P, Robberecht P. Fragments of pituitary adenylate cyclase activating polypeptide discriminate between type I and II recombinant receptors. *Eur J Pharmacol* 1995;287(1):7-11.
- [26] Guo S, Vollesen AL, Hansen RD, Esserlind AL, Amin FM, Christensen AF, Olesen J, Ashina M. Part I: Pituitary adenylate cyclase-activating polypeptide-38 induced migraine-like attacks in patients with and without familial aggregation of migraine. *Cephalalgia* 2017;37:125-135.
- [27] Harmar AJ, Fahrenkrug J, Gozes I, Laburthe M, May V, Pisegna JR, Vaudry D, Vaudry H, Waschek JA, Said SI. IUPHAR Reviews 1: Pharmacology and functions of receptors for vasoactive intestinal peptide and pituitary adenylate cyclase-activating polypeptide. *Br J Pharmacol* 2012;166:4-17.
- [28] Hashimoto H, Nogi H, Mori K, Ohishi H, Shigemoto R, Yamamoto K, Matsuda T, Mizuno N, Nagata S, Baba A. Distribution of the mRNA for a pituitary adenylate cyclase-activating polypeptide receptor in the rat brain: an in situ hybridization study. *J Comp Neurol* 1996;371(4):567-577.
- [29] Hewitt DJ, Aurora SK, Dodick DW, Goadsby PJ, Ge J, Bachman R, Taraborelli D, Fan X, Assaid C, Lines C, Ho TW. Randomized controlled trial of the CGRP receptor

antagonist, MK-3207, in the acute treatment of migraine. Cephalalgia 2011;31:712-722.

[30] Ho AP, Dahlof CG, Silberstein SD, Saper JR, Ashina M, Kost JT, Froman S, Leibensperger H, Lines CR, Ho TW. Randomized, controlled trial of telcagepant over four migraine attacks. Cephalalgia 2010;30(12):1443-1457.

[31] Ho TW, Ferrari MD, Dodick DW, Galet V, Kost J, Fan X, Leibensperger H, Froman S, Assaid C, Lines C, Koppen H, Winner PK. Efficacy and tolerability of MK-0974 (telcagepant), a new oral antagonist of calcitonin gene-related peptide receptor, compared with zolmitriptan for acute migraine: a randomised, placebo-controlled, parallel-treatment trial. Lancet 2008;372:2115-2123.

[32] Hoffmann J, Martins-Oliveira M, Akerman S, Suprongsinchai W, Xu C, Goadsby PJ. PAC-1 receptor antibody modulates nociceptive trigeminal activity in rat. Cephalalgia 2016;36(1S):141.

[33] Hoffmann J, Martins-Oliveira M, Akerman S, Suprongsinchai W, Xu C, Goadsby PJ. Nociceptive trigeminal neurotransmission is inhibited by a PAC-1 receptor antibody in an in vivo model relevant to migraine. Cephalalgia 2017;31(1S):3-4.

[34] Hoffmann J, May A. Diagnosis, pathophysiology, and management of cluster headache. Lancet Neurol 2018;17(1):75-83.

[35] Jansen-Olesen I, Baun M, Amrutkar DV, Ramachandran R, Christophersen DV, Olesen J.

PACAP-38 but not VIP induces release of CGRP from trigeminal nucleus caudalis via a receptor distinct from the PAC1 receptor. *Neuropeptides* 2014;48(2):53-64.

[36] Laburthe M, Couvineau A, Tan V. Class II G protein-coupled receptors for VIP and

PACAP: structure, models of activation and pharmacology. *Peptides*

2007;28(9):1631-1639.

[37] Lassen LH, Haderslev PA, Jacobsen VB, Iversen HK, Sperling B, Olesen J. CGRP may play a

causative role in migraine. *Cephalalgia* 2002;22:54-61.

[38] Marcus R, Goadsby PJ, Dodick D, Stock D, Manos G, Fischer TZ. BMS-927711 for the

Acute Treatment of Migraine: A Double-Blind, Randomized, Placebo Controlled,

Dose-ranging Trial. *Cephalalgia* 2014;34:114-125.

[39] Martinez JM, Goadsby PJ, Dodick DW, Bardos JN, Oakes TMM, Millen BA, Zhou C,

Dowsett SA, Aurora S, Yang JY, Conley RR. Study CGAL: A placebo-controlled study of

galcanezumabin patients with episodic cluster headache: Results from the 8-week

double-blind treatment phase. *Cephalalgia* 2018;38(1S):145-146.

[40] Miyata A, Arimura A, Dahl RR, Minamino N, Uehara A, Jiang L, Culler MD, Coy DH.

Isolation of a novel 38 residue-hypothalamic polypeptide which stimulates adenylate cyclase in pituitary cells. *Biochem Biophys Res Commun* 1989;164(1):567-574.

- [41] Moro O, Lerner EA. Maxadilan, the vasodilator from sand flies, is a specific pituitary adenylate cyclase activating peptide type I receptor agonist. *J Biol Chem* 1997;272(2):966-970.
- [42] Olesen J, Diener HC, Husstedt IW, Goadsby PJ, Hall D, Meier U, Pollentier S, Lesko LM. Calcitonin gene-related peptide receptor antagonist BIBN 4096 BS for the acute treatment of migraine. *New England Journal of Medicine* 2004;350:1104-1110.
- [43] Rahmann A, Wienecke T, Hansen JM, Fahrenkrug J, Olesen J, Ashina M. Vasoactive intestinal peptide causes marked cephalic vasodilatation but does not induce migraine. *Cephalalgia* 2008;28:226-236.
- [44] Schytz HW, Birk S, Wienecke T, Kruuse C, Olesen J, Ashina M. PACAP38 induces migraine-like attacks in patients with migraine without aura. *Brain* 2009;132:16-25.
- [45] Schytz HW, Birk S, Wienecke T, Kruuse C, Olesen J, Ashina M. PACAP38 induces migraine-like attacks in patients with migraine without aura. *Brain* 2009;132(Pt 1):16-25.
- [46] Silberstein SD, Aycardi E, Bigal ME, Blankenbiller T, Dodick DW, Goadsby PJ, Grozinski-Wolff M, Ma M, Yang R, Yeung P. Fremanezumab for chronic migraine preventive treatment. *New England Journal of Medicine* 2017;377:2113-2122.

- [47] Silberstein SD, Dodick DW, Bigal ME, Yeung PP, Goadsby PJ, Blankenbiller T, Grozinski-Wolff M, Yang R, Ma Y, Aycardi E. Fremanezumab for the Preventive Treatment of Chronic Migraine. *N Engl J Med* 2017;377(22):2113-2122.
- [48] Skljarevski V, Oakes TM, Zhang Q, Ferguson MB, Martinez J, Camporeale A, Johnson KW, Shan Q, Carter J, Schacht A, Goadsby PJ, Dodick DW. Galcanezumab for Episodic Migraine Prevention: A Randomized Phase 2b Placebo-controlled Dose-ranging Clinical Trial. *JAMA Neurology* 2018;75:187-193.
- [49] Spengler D, Waeber C, Pantaloni C, Holsboer F, Bockaert J, Seeburg PH, Journot L. Differential signal transduction by five splice variants of the PACAP receptor. *Nature* 1993;365(6442):170-175.
- [50] Storer RJ, Akerman S, Goadsby PJ. Calcitonin gene-related peptide (CGRP) modulates nociceptive trigeminovascular transmission in the cat. *British Journal of Pharmacology* 2004;142:1171-1181.
- [51] Tuka B, Helyes Z, Markovics A, Bagoly T, Szolcsanyi J, Szabo N, Toth E, Kincses ZT, Vecsei L, Tajti J. Alterations in PACAP-38-like immunoreactivity in the plasma during ictal and interictal periods of migraine patients. *Cephalalgia* 2013;33:1085-1095.
- [52] Tuka B, Szabo N, Toth E, Kincses ZT, Pardutz A, Szok D, Kortesi T, Bagoly T, Helyes Z, Edvinsson L, Vecsei L, Tajti J. Release of PACAP-38 in episodic cluster headache patients - an exploratory study. *J Headache Pain* 2016;17(1):69.

[53] Vos T, Abajobir AA, Abate KH, Abbafati C, Abbas KM, Abd-Allah F, Abdulkader RS, Abdulle AM, Abebo TA, Abera SF, Aboyans V, Abu-Raddad LJ, Ackerman IN, Adamu AA, Adetokunboh O, Afarideh M, Afshin A, Agarwal SK, Aggarwal R, Agrawal A, Agrawal S, Ahmadieh H, Ahmed MB, Aichour MTE, Aichour AN, Aichour I, Aiyar S, Akinyemi RO, Akseer N, Al Lami FH, Alahdab F, Al-Aly Z, Alam K, Alam N, Alam T, Alasfoor D, Alene KA, Ali R, Alizadeh-Navaei R, Alkerwi Aa, Alla F, Allebeck P, Allen C, Al-Maskari F, Al-Raddadi R, Alsharif U, Alsowaidi S, Altirkawi KA, Amare AT, Amini E, Ammar W, Amoako YA, Andersen HH, Antonio CAT, Anwari P, Ärnlöv J, Artaman A, Aryal KK, Asayesh H, Asgedom SW, Assadi R, Atey TM, Atnafu NT, Atre SR, Avila-Burgos L, Avokphako EFGA, Awasthi A, Bacha U, Badawi A, Balakrishnan K, Banerjee A, Bannick MS, Barac A, Barber RM, Barker-Collo SL, Bärnighausen T, Barquera S, Barregard L, Barrero LH, Basu S, Battista B, Battle KE, Baune BT, Bazargan-Hejazi S, Beardsley J, Bedi N, Beghi E, Béjot Y, Bekele BB, Bell ML, Bennett DA, Bensenor IM, Benson J, Berhane A, Berhe DF, Bernabé E, Betsu BD, Beuran M, Beyene AS, Bhala N, Bhansali A, Bhatt S, Bhutta ZA, Biadgilign S, Bicer BK, Bienhoff K, Bikbov B, Birungi C, Biryukov S, Bisanzio D, Bizuayehu HM, Boneya DJ, Boufous S, Bourne RRA, Brazinova A, Brugha TS, Buchbinder R, Bulto LNB, Bumgarner BR, Butt ZA, Cahuana-Hurtado L, Cameron E, Car M, Carabin H, Carapetis JR, Cárdenas R, Carpenter DO, Carrero JJ, Carter A, Carvalho F, Casey DC, Caso V, Castañeda-Orjuela CA, Castle CD, Catalá-López F, Chang H-Y, Chang J-C, Charlson FJ, Chen H, Chibalabala M, Chibueze CE, Chisumpa VH, Chitheer AA, Christopher DJ, Ciobanu LG, Cirillo M, Colombara D, Cooper C, Cortesi PA, Criqui MH, Crump JA, Dadi AF, Dalal K, Dandona L, Dandona R, das Neves J, Davitoiu DV, de Courten B, De Leo DD, Defo BK, Degenhardt L, Deiparine S, Dellavalle RP, Deribe K, Des Jarlais DC, Dey S, Dharmaratne SD, Dhillon PK, Dicker

D, Ding EL, Djalalinia S, Do HP, Dorsey ER, dos Santos KPB, Douwes-Schultz D, Doyle
 KE, Driscoll TR, Dubey M, Duncan BB, El-Khatib ZZ, Ellerstrand J, Enayati A, Endries
 AY, Ermakov SP, Erskine HE, Eshrati B, Eskandarieh S, Esteghamati A, Estep K, Fanuel
 FBB, Farinha CSES, Faro A, Farzadfar F, Fazeli MS, Feigin VL, Fereshtehnejad S-M,
 Fernandes JC, Ferrari AJ, Feyissa TR, Filip I, Fischer F, Fitzmaurice C, Flaxman AD, Flor
 LS, Foigt N, Foreman KJ, Franklin RC, Fullman N, Fürst T, Furtado JM, Futran ND,
 Gakidou E, Ganji M, Garcia-Basteiro AL, Gebre T, Gebrehiwot TT, Geleto A, Gemechu
 BL, Gesesew HA, Gething PW, Ghajar A, Gibney KB, Gill PS, Gillum RF, Ginawi IAM,
 Giref AZ, Gishu MD, Giussani G, Godwin WW, Gold AL, Goldberg EM, Gona PN,
 Goodridge A, Gopalani SV, Goto A, Goulart AC, Griswold M, Gughani HC, Gupta R,
 Gupta R, Gupta T, Gupta V, Hafezi-Nejad N, Hailu GB, Hailu AD, Hamadeh RR, Hamidi
 S, Handal AJ, Hankey GJ, Hanson SW, Hao Y, Harb HL, Hareri HA, Haro JM, Harvey J,
 Hassanvand MS, Havmoeller R, Hawley C, Hay SI, Hay RJ, Henry NJ, Heredia-Pi IB,
 Hernandez JM, Heydarpour P, Hoek HW, Hoffman HJ, Horita N, Hosgood HD, Hostiuc
 S, Hotez PJ, Hoy DG, Htet AS, Hu G, Huang H, Huynh C, Iburg KM, Igumbor EU, Ikeda
 C, Irvine CMS, Jacobsen KH, Jahanmehr N, Jakovljevic MB, Jassal SK, Javanbakht M,
 Jayaraman SP, Jeemon P, Jensen PN, Jha V, Jiang G, John D, Johnson SC, Johnson CO,
 Jonas JB, Jürisson M, Kabir Z, Kadel R, Kahsay A, Kamal R, Kan H, Karam NE, Karch A,
 Karema CK, Kasaeian A, Kassa GM, Kassaw NA, Kassebaum NJ, Kastor A, Katikireddi
 SV, Kaul A, Kawakami N, Keiyoro PN, Kengne AP, Keren A, Khader YS, Khalil IA, Khan
 EA, Khang Y-H, Khosravi A, Khubchandani J, Kiadaliri AA, Kieling C, Kim YJ, Kim D, Kim
 P, Kimokoti RW, Kinfu Y, Kisa A, Kissimova-Skarbek KA, Kivimaki M, Knudsen AK,
 Kokubo Y, Kolte D, Kopec JA, Kosen S, Koul PA, Koyanagi A, Kravchenko M,
 Krishnaswami S, Krohn KJ, Kumar GA, Kumar P, Kumar S, Kyu HH, Lal DK, Lalloo R,

Lambert N, Lan Q, Larsson A, Lavados PM, Leasher JL, Lee PH, Lee J-T, Leigh J,
 Leshargie CT, Leung J, Leung R, Levi M, Li Y, Li Y, Li Kappe D, Liang X, Liben ML, Lim
 SS, Linn S, Liu PY, Liu A, Liu S, Liu Y, Lodha R, Logroscino G, London SJ, Looker KJ,
 Lopez AD, Lorkowski S, Lotufo PA, Low N, Lozano R, Lucas TCD, Macarayan ERK,
 Magdy Abd El Razek H, Magdy Abd El Razek M, Mahdavi M, Majdan M, Majdzadeh R,
 Majeed A, Malekzadeh R, Malhotra R, Malta DC, Mamun AA, Manguerra H,
 Manhertz T, Mantilla A, Mantovani LG, Mapoma CC, Marczak LB, Martinez-Raga J,
 Martins-Melo FR, Martopullo I, März W, Mathur MR, Mazidi M, McAlinden C,
 McGaughey M, McGrath JJ, McKee M, McNellan C, Mehata S, Mehndiratta MM,
 Mekonnen TC, Memiah P, Memish ZA, Mendoza W, Mengistie MA, Mengistu DT,
 Mensah GA, Meretoja TJ, Meretoja A, Mezgebe HB, Micha R, Millear A, Miller TR,
 Mills EJ, Mirarefin M, Mirrakhimov EM, Misganaw A, Mishra SR, Mitchell PB,
 Mohammad KA, Mohammadi A, Mohammed KE, Mohammed S, Mohanty SK,
 Mokdad AH, Mollenkopf SK, Monasta L, Montico M, Moradi-Lakeh M, Moraga P,
 Mori R, Morozoff C, Morrison SD, Moses M, Mountjoy-Venning C, Mruts KB, Mueller
 UO, Muller K, Murdoch ME, Murthy GVS, Musa KI, Nachega JB, Nagel G, Naghavi M,
 Naheed A, Naidoo KS, Naldi L, Nangia V, Natarajan G, Negasa DE, Negoï RI, Negoï I,
 Newton CR, Ngunjiri JW, Nguyen TH, Nguyen QL, Nguyen CT, Nguyen G, Nguyen M,
 Nichols E, Ningrum DNA, Nolte S, Nong VM, Norrving B, Noubiap JJN, O'Donnell MJ,
 Ogbo FA, Oh I-H, Okoro A, Oladimeji O, Olagunju TO, Olagunju AT, Olsen HE,
 Olusanya BO, Olusanya JO, Ong K, Opio JN, Oren E, Ortiz A, Osgood-Zimmerman A,
 Osman M, Owolabi MO, Pa M, Pacella RE, Pana A, Panda BK, Papachristou C, Park E-
 K, Parry CD, Parsaeian M, Patten SB, Patton GC, Paulson K, Pearce N, Pereira DM,
 Perico N, Pesudovs K, Peterson CB, Petzold M, Phillips MR, Pigott DM, Pillay JD, Pinho

C, Plass D, Pletcher MA, Popova S, Poulton RG, Pourmalek F, Prabhakaran D, Prasad
 NM, Prasad N, Purcell C, Qorbani M, Quansah R, Quintanilla BPA, Rabiee RHS, Radfar
 A, Rafay A, Rahimi K, Rahimi-Movaghar A, Rahimi-Movaghar V, Rahman MHU,
 Rahman M, Rai RK, Rajsic S, Ram U, Ranabhat CL, Rankin Z, Rao PC, Rao PV, Rawaf S,
 Ray SE, Reiner RC, Reinig N, Reitsma MB, Remuzzi G, Renzaho AMN, Resnikoff S,
 Rezaei S, Ribeiro AL, Ronfani L, Roshandel G, Roth GA, Roy A, Rubagotti E, Ruhago
 GM, Saadat S, Sadat N, Safdarian M, Safi S, Safiri S, Sagar R, Sahathevan R, Salama J,
 Saleem HOB, Salomon JA, Salvi SS, Samy AM, Sanabria JR, Santomauro D, Santos IS,
 Santos JV, Santric Milicevic MM, Sartorius B, Satpathy M, Sawhney M, Saxena S,
 Schmidt MI, Schneider IJC, Schöttker B, Schwebel DC, Schwendicke F, Seedat S,
 Sepanlou SG, Servan-Mori EE, Setegn T, Shackelford KA, Shaheen A, Shaikh MA,
 Shamsipour M, Shariful Islam SM, Sharma J, Sharma R, She J, Shi P, Shields C, Shifa
 GT, Shigematsu M, Shinohara Y, Shiri R, Shirkoohi R, Shirude S, Shishani K, Shrima
 MG, Sibai AM, Sigfusdottir ID, Silva DAS, Silva JP, Silveira DGA, Singh JA, Singh NP,
 Sinha DN, Skiadaresi E, Skirbekk V, Slepak EL, Sligar A, Smith DL, Smith M, Sobaih
 BHA, Sobngwi E, Sorensen RJD, Sousa TCM, Sposato LA, Sreeramareddy CT,
 Srinivasan V, Stanaway JD, Stathopoulou V, Steel N, Stein MB, Stein DJ, Steiner TJ,
 Steiner C, Steinke S, Stokes MA, Stovner LJ, Strub B, Subart M, Sufiyan MB, Sunguya
 BF, Sur PJ, Swaminathan S, Sykes BL, Sylte DO, Tabarés-Seisdedos R, Taffere GR,
 Takala JS, Tandon N, Tavakkoli M, Taveira N, Taylor HR, Tehrani-Banihashemi A,
 Tekelab T, Terkawi AS, Tesfaye DJ, Tessema B, Thamsuwan O, Thomas KE, Thrift AG,
 Tiruye TY, Tobe-Gai R, Tollanes MC, Tonelli M, Topor-Madry R, Tortajada M, Touvier
 M, Tran BX, Tripathi S, Troeger C, Truelsen T, Tsoi D, Tuem KB, Tuzcu EM, Tyrovolas S,
 Ukwaja KN, Undurraga EA, Uneke CJ, Updike R, Uthman OA, Uzochukwu BSC, van

Boven JFM, Varughese S, Vasankari T, Venkatesh S, Venketasubramanian N,
 Vidavalur R, Violante FS, Vladimirov SK, Vlassov VV, Vollset SE, Wadilo F, Wakayo T,
 Wang Y-P, Weaver M, Weichenthal S, Weiderpass E, Weintraub RG, Werdecker A,
 Westerman R, Whiteford HA, Wijeratne T, Wiysonge CS, Wolfe CDA, Woodbrook R,
 Woolf AD, Workicho A, Xavier D, Xu G, Yadgir S, Yaghoubi M, Yakob B, Yan LL, Yano Y,
 Ye P, Yimam HH, Yip P, Yonemoto N, Yoon S-J, Yotebieng M, Younis MZ, Zaidi Z, Zaki
 MES, Zegeye EA, Zenebe ZM, Zhang X, Zhou M, Zipkin B, Zodpey S, Zuhlke LJ, Murray
 CJL. Global, regional, and national incidence, prevalence, and years lived with
 disability for 328 diseases and injuries for 195 countries, 1990-2016: a systematic
 analysis for the Global Burden of Disease Study 2016. *The Lancet*
 2017;390(10100):1211-1259.

[54] Voss T, Lipton RB, Dodick DW, Dupre N, Ge JY, Bachman R, Assaid C, Aurora SK,

Michelson D. A phase IIb randomized, double-blind, placebo-controlled trial of
 ubrogepant for the acute treatment of migraine. *Cephalalgia* 2016;36:887-898.

[55] Zagami AS, Edvinsson L, Goadsby PJ. Pituitary adenylate cyclase activating polypeptide
 and Migraine. *Annals of Clinical and Translational Neurology* 2014;1:1036-1040.

[56] Zagami AS, Edvinsson L, Hoskin KL, Goadsby PJ. Stimulation of the superior sagittal sinus
 causes extracranial release of PACAP. *Cephalalgia* 1995;15(Suppl 14):109.

[57] Zagami AS, Goadsby PJ, Edvinsson L. Stimulation of the superior sagittal sinus in the cat
 causes release of vasoactive peptides. *Neuropeptides* 1990;16:69-75.

[58] Zhang Y, Danielsen N, Sundler F, Mulder H. Pituitary adenylate cyclase-activating peptide is upregulated in sensory neurons by inflammation. *Neuroreport* 1998;9(12):2833-2836.

Figures

Fig. 1: Antagonist activity of Ab181 at rat rPAC1 CHO cell (rPAC1-CHO) against PACAP38 or maxadilan

The antagonist activity of Ab181 at the rat PAC₁ receptor was carried out by measuring the potency of Ab181 in inhibiting EC₈₀ of agonist PACAP38 or maxadilan stimulated cAMP production. PACAP6-38, a PAC₁ receptor antagonist, was used as a positive control. Full antagonist activity of Ab181 against both PACAP38 (A) and maxadilan (B) induced cAMP accumulation was observed in rPAC1-CHO cells with IC₅₀ of 20 ± 3.3 nM ($n = 3$) and 4.5 ± 0.1 nM ($n = 2$) against respective agonist.

Fig 2.1: Prophylactic effect of Ab181 on maxadilan-induced increase in DBF in rats 48 h post drug treatment

Rats were given i.v. bolus injection of Ab181 (0.1, 0.3, 1 and 10 mg/kg, $n = 6-7$ rats/group). 48 h later, after obtaining a post-drug baseline (BL), maxadilan at a dose of 10 ng (in 20 μ l DPBS) was injected intradermally. The dermal blood flow (DBF) was measured by a laser Doppler imager. Graph A: The increase in DBF (change from BL) caused by maxadilan injection over 60 min was expressed as % change from BL [= 100 x (individual post-maxadilan flux - individual BL flux)/individual BL flux]. Graph B: % change from BL (left-axis) versus resulting serum concentrations (right-axis) at 15 min post maxadilan; Solid triangle represents % change from BL at 48 h post drug treatment (left-axis, mean \pm SEM), open bar represents serum concentration at 48 h post drug treatment (right-axis, mean \pm SD). *** $p < 0.001$, **** $p < 0.0001$ compared to vehicle-treated group by one-way ANOVA followed by Dunnett's MCT.

Fig 2.2: Effect of Ab181 on a maxadilan-induced increase in DBF in rats 3.25 h post treatment

Rats were given i.v. bolus injection of Ab181 (0.1, 0.3, 1, 3 and 10 mg/kg, n = 7-8 rats/group). 3 h later, after obtaining a post-drug baseline (BL), maxadilan at a dose of 10 ng (in 20 µl DPBS) was injected intradermally. The dermal blood flow was measured by a laser Doppler imager. Solid triangle represents % change from baseline (mean ± SEM, left axis), open bar represents resulting serum concentration (mean ± SD, right axis). Control: Dummy antibody + vehicle (n = 13 rats). **p<0.01, ****p<0.0001 compared to control group by one-way ANOVA followed by Dunnett's MCT.

Fig 2.3: Time-course of effect of Ab181 (at 10 mg/kg) on a maxadilan-induced increase in DBF in rats

Rats were given i.v. bolus injection of Ab181 at 10 mg/kg with pretreatment time at 0.5, 3, 6.25, 48 and 168 h prior to maxadilan challenge (n = 3-8 rats/group). After obtaining a post-drug baseline (BL), maxadilan at a dose of 10 ng (in 20 µl DPBS) was injected intradermally. The dermal blood flow was measured by a laser Doppler imager. Solid triangle represents % Inhibition (mean ± SEM, left-axis), open bar represents resulting serum concentration (mean ± SD, right-axis).

Figure 3.1: Timeline of the electrophysiological *in vivo* experiments

Following the surgical procedure and the recording of baseline activity either Ab181 (10 mg/kg) or vehicle (A5Su) were injected intravenously. After a resting period of 2.5 h to allow the distribution of the antibody throughout the circulation and the relevant tissues, sumatriptan or its vehicle were administered intravenously. Following another resting

period of 30 min to allow sumatriptan to act, stimulus-evoked and spontaneous neuronal activity was recorded in the TCC.

Figure 3.2: Effect of Ab181 on neuronal activity in the TCC

Ab181 effectively inhibited stimulus-evoked neuronal activity in the TCC (A). The maximum inhibition was very similar to the one observed after the intravenous administration of sumatriptan (B). In contrast, an inhibiting effect on spontaneous background activity was not observed (C). Ab181 did not affect arterial blood pressure throughout the entire course of the experiment (D).

Figure 4: Distribution of Ab181 in tissues after intravenous dosing

Ex vivo detection of the IgG-portion of Ab181 or control antibody after i.v. dosing in dural vessels, trigeminal ganglion (TG) and sphenopalatine ganglion (SPG) as well as spinal trigeminal nucleus caudalis (TNC), superior salivatory nucleus (SSN) and thalamus. Staining of neuronal and glial structures is detected in dura, TG and SPG of Ab181-dosed rats, but not in TNC, SSN or thalamus. Only background staining is detected after intravenous dosing with control antibody (lower panels). Scale bars 50 μ m.

Table 1: Antagonist Activities (IC₅₀) of Ab181

The antagonist activity of Ab181 at the rat PAC₁ receptor (rPAC₁) was carried out by measuring the potency of Ab181 in inhibiting EC₈₀ of agonist PACAP38 or maxadilan stimulated cAMP production. PACAP6-38, a PAC₁ receptor antagonist, was used as a positive control.

Similar assay was used to determine the activity of Ab181 at rat VPAC₁ (rVPAC₁) and rat VPAC₂ (rVPAC₂) receptors against the EC₈₀ of agonist VIP.


	Agonist	rPAC ₁		rVPAC ₁	rVPAC ₂
		PACAP38	maxadilan	VIP	VIP
Ab181 (IC ₅₀ , nM)		20 ± 3.3 (n = 3)	4.5 ± 0.1 (n=2)	>1000 (n=2)	>1000 (n=2)
PACAP6-38 (IC ₅₀ , nM)		266 ± 77 (n = 3)	2.9 ± 1.5 (n=2)	430 ± 79 (n=2)	38 ± 4 (n=2)

IC₅₀ values from these experiments (Table 1) are expressed as mean ± SD, and “n” represents the number of separate experiments.


1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Summary


PAC₁ receptor blockade using a monoclonal antibody inhibits nociceptive neuronal activity in the trigeminocervical complex.




A



B









Figure 3.1







Figure 3.2

[Click here to access/download;Figure;Figure 3-2 new.tif](#)





Open Access License Agreement

This OPEN ACCESS LICENSE AGREEMENT (this "Agreement"), dated as of.

DATE

(the "Effective Date"), by and between Wolters Kluwer Health, Inc., operating as Medical Research / Lippincott Williams & Wilkins, a Delaware corporation, having its principal place of business at Two Commerce Square, 2001 Market Street, Philadelphia, PA 19103 (the "Publisher"), and the corresponding author listed on Schedule A to this Agreement (the "Author", and together with the Publisher, the "Parties").

1. Grant of License

The Author hereby grants to the Publisher and its Affiliates the exclusive, worldwide, royalty free, perpetual (for the duration of the applicable copyright) right and license to use the Work for all commercial or educational purposes, including, but not limited to, publishing, reproducing, marketing, distributing (themselves and through distributors), sublicensing, and selling copies of the Work throughout the world for the Term. If the Author is a United States government employee, such license grant shall be limited to the extent the Author is able to grant such license.

2. Warranties, Indemnification, and Limitation of Liability

a. The Author represents and warrants that:

(i) it has the right and power to enter into this Agreement, to grant the rights and licenses granted pursuant to this Agreement, and to perform all of its other obligations contained in this Agreement;

(ii) it has not previously assigned, transferred or otherwise encumbered the rights or licenses granted

pursuant to this Agreement; and that the person executing this Agreement on the Author's behalf is authorized to do so;

(iii) the Work and the licenses granted herein do not and will not infringe upon, violate or misappropriate any intellectual property rights or any other proprietary right, contract or other right or interest of any third party;

(iv) if the Work is a multi-authored Work, the Author has obtained written permission from each author of the Work to enter into this Agreement on behalf such author, and each such author has read, understands and has agreed to the terms of this Agreement; and

(v) the Author has obtained any necessary releases and permissions to quote from other sources in the Work and to include any works and materials in the Work and all such releases and permissions are in full force and effect.

b. The Author hereby indemnifies the Publisher and its directors, officers, employees, agents, and representatives and agrees to defend and hold them harmless from and against any and all liability, damage, loss, costs or expenses (including reasonable attorney's fees and costs of settlement) incurred by any such party arising out of, or relating to any misrepresentation in, or breach or alleged breach of the Author's representations or warranties in this Agreement. If the Author fails to promptly or diligently pursue any defense of any indemnified party, the indemnified parties, or any of them, may assume such defense at the Author's expense. The obligations of this indemnification will survive any termination or expiration of this Agreement.

c. The Publisher represents and warrants that it has the right and power to enter into this Agreement and to perform its obligations contained in this Agreement, and that the person executing this Agreement on the Publisher's behalf is authorized to do so.

d. The Publisher hereby indemnifies the Author and agrees to defend and hold the Author harmless from and against any and all liability, damage, loss, costs or expenses (including reasonable attorney's fees and costs of settlement) incurred by the Author arising out of, or relating to any misrepresentation in, or breach or alleged breach of the Publisher's representations or warranties in this Agreement. If the Publisher fails to promptly or diligently pursue any defense of the Author, the Author may assume such defense at the Publisher's expense. The obligations of this indemnification will survive any termination or expiration of this Agreement.

e. EXCEPT AS OTHERWISE SET FORTH IN THIS AGREEMENT, NEITHER PARTY MAKES ANY OTHER, AND HEREBY DISCLAIMS ALL OTHER, REPRESENTATIONS AND WARRANTIES OF ANY KIND, WHETHER EXPRESS, IMPLIED, STATUTORY OR OTHERWISE, INCLUDING, WITHOUT LIMITATION, WARRANTIES OF TITLE, MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, NONINFRINGEMENT, OR THE ABSENCE OF LATENT OR OTHER DEFECTS, ACCURACY, OR THE PRESENCE OR ABSENCE OF ERRORS, WHETHER OR NOT DISCOVERABLE.

f. EXCEPT TO THE EXTENT REQUIRED BY APPLICABLE LAW, IN NO EVENT WILL EITHER PARTY BE LIABLE TO THE OTHER PARTY BASED UPON ANY LEGAL THEORY FOR ANY SPECIAL, INCIDENTAL, CONSEQUENTIAL, PUNITIVE, OR EXEMPLARY DAMAGES ARISING OUT OF THIS LICENSE OR THE USE OF THE WORK, EVEN IF A PARTY HAS BEEN ADVISED OF THE POSSIBILITY OF SUCH DAMAGES.

3. Creative Commons License.

Creative Commons Licenses are subject to items selected in item 1, 2 and 3 in the Schedule B.

a. CCBY-NC-ND – NonCommercial-NonDerivatives Creative Commons License

The Author acknowledges and agrees that the Work will be published by the Publisher in (the "Journal") and made freely available to users under the terms of the Attribution-NonCommercial-NoDerivs 4.0 Creative Commons License, as currently displayed at <http://creativecommons.org/licenses/by-nc-nd/4.0/legalcode> (the "CC BY-NC-ND"). The Author acknowledges and agrees

that that Publisher is the exclusive "Licensor", as defined in the CC BY-NC-ND, of the Work and that the Publisher may make the Work freely available to all users under the terms of the CC BY-NC-ND.

b. CCBY – Creative Commons License

The Author acknowledges and agrees that the Work will be published by the Publisher in (the "Journal") and made freely available to users under the terms of the Attribution 4.0 Creative Commons License, as currently displayed at <http://creativecommons.org/licenses/by/4.0/legalcode> (the "CC BY"). The Author acknowledges and agrees that that Publisher is the exclusive "Licensor", as defined in the CC BY, of the Work and that the Publisher may make the Work freely available to all users under the terms of the CC BY.

c. CCBY-NC – NonCommercial Creative Commons License

The Author acknowledges and agrees that the Work will be published by the Publisher in (the "Journal") and made freely available to users under the terms of the Attribution-NonCommercial 4.0 Creative Commons License, as currently displayed at <http://creativecommons.org/licenses/by-nc/4.0/legalcode> (the "CC BY-NC"). The Author acknowledges and agrees that that Publisher is the exclusive "Licensor", as defined in the CC BY-NC, of the Work and that the Publisher may make the Work freely available to all users under the terms of the CC BY-NC.

4. Royalties.

The Author acknowledges and agrees that this Agreement entitles the Author to no royalties or fees. To the maximum extent permitted by law, the Author waives any and all rights the Author may have to collect royalties or other fees in relation to the Work or in respect of any use of the Work by the Publisher or its sublicensees.

5. Miscellaneous.

a. Assignment. This Agreement may not be assigned or transferred, in whole or in part, by either party without the prior written consent of the other party. Notwithstanding the above, the Publisher may assign this Agreement without the written consent of the Author (i) to an entity succeeding, whether by sale, merger or other corporate reorganization,

to substantially all of the Publisher's assets and business activity, or (ii) to a corporation or organization that obtains the right to publish the Journal from the Publisher. The Publisher may assign this Agreement to any of its affiliates. This Agreement will be binding upon and inure to the benefit of the parties hereto and their respective successors and permitted assigns.

b. Counterparts. This Agreement may be executed in two or more counterparts, each of which shall be deemed an original, but all of which together shall constitute one and the same document. Facsimile or Portable Document Format (PDF) signatures will be deemed original signatures for purposes of this Agreement.

c. Entire Agreement; Amendment. This Agreement sets forth the entire agreement of the parties on the subject hereof and supersedes all previous or contemporaneous oral or written representations or agreements relating to the rights and duties provided herein, and may not be modified or amended except by written agreement of the parties.

d. Force Majeure. Neither party shall be liable for any default or delay on its part in performing any obligation under this Agreement if such default or delay is caused by natural disaster, accident, war, civil disorder, strike or any other cause beyond the reasonable control of such party. In the event that either party is prevented by such an occurrence or circumstance for a period of more than ninety (90) days from fulfilling its obligations under this Agreement, the other party may terminate this Agreement upon thirty (30) days' written notice.

e. Governing Law. This Agreement shall be governed in all respects according to the laws of the State of New York without giving effect to the principles of conflict of law thereof.

f. Headings. All headings are for reference purposes only and shall not affect the meaning or interpretation of any provision hereof.

g. Severability. If any provision of this Agreement is held to be illegal, invalid, or unenforceable under the present or future laws, then such provision shall be revised by a court of competent jurisdiction to be enforceable if permitted under applicable law, and otherwise shall be fully severable. In any event, this Agreement shall be construed and enforced as if such illegal, invalid, or unenforceable provision had never comprised a part of this Agreement, and the remaining provisions of this Agreement shall remain in full force and effect and shall not be affected by the illegal, invalid, or unenforceable provision or by its severance from this Agreement.

h. Status of the Parties. The parties are independent contractors. Nothing in this Agreement is intended to or shall be construed to constitute or establish any agency, joint venture, partnership or fiduciary relationship between the parties, and neither party has the right or authority to bind the other party nor shall either party be responsible for the acts or omissions of the other.

i. Waiver; Amendment. The waiver by either party of or the failure by either party to claim a breach of any provision of this Agreement shall not be, or be held to be, a waiver of any subsequent breach or affect in any way the further effectiveness of any such provision. No term or condition of this Agreement may be waived except by an agreement by the parties in writing.

j. Waiver of Jury Trial. EACH PARTY HEREBY WAIVES ITS RIGHT TO A JURY TRIAL IN CONNECTION WITH ANY DISPUTE OR LEGAL PROCEEDING ARISING OUT OF THIS AGREEMENT OR THE SUBJECT MATTER HEREOF.

[Signature Page Follows]

Schedule A

This Schedule A must be completed by Author in its entirety. The Publisher is unable to publish the Work unless this Schedule A is completely filled out.

Article Tracking #

Article Title (the “Work”)

Corresponding Author Name (the “Author”)

Copyright Owner’s Name

Name of Journal in which Work is to be Published

Schedule B

This Schedule B must be completed by Author in its entirety. The Publisher is unable to publish the Work unless this Schedule B is completely filled out.

MANDATED FUNDING POLICY DISCLOSURE

1. Choose a funder from the drop down list. If any of the following are selected please complete Item 2.

NOTE: If you are a World Health Organization Employee and are required to publish under the Creative Commons CCBY IGO license, then do not complete this form. Instead, please contact the Editorial Office for the separate WHO Employee license agreement.

2. If you have selected funding from the above list in 1., please disclose the Open Access option to which the Work will be subject. Selecting “Gold Route” will ensure that your work is published under the Creative Commons CCBY license.

☐ Gold route

☐ Green route

NOTE: If the “Gold” route has been selected, Section 3.b. of the Agreement will apply to the Work, and neither Section 3.a. nor Section 3.c. of the Agreement will apply to the Work. If the “Green” route has been selected, Section 3.c. of the Agreement will apply to the Work after an embargo, and neither Section 3.a. nor Section 3.b. of the Agreement will apply to the Work.

3. ☐ This Schedule B is inapplicable to the Work.

NOTE: If author has selected Item 3, Section 3.a. on the Agreement will apply to the Work, and neither Section 3.b. nor Section 3.c. of the Agreement will apply to the Work.

GOVERNMENT EMPLOYEES

4. ☐ This work was created in the course of an author's employment by the United States Government

If the Work or a portion of it has been created in the course of any author's employment by the United States Government, check the "Government" box at the end of this form. A work prepared by a government employee as part of his or her official duties is called a "work of the U.S. Government" and is not subject to copyright. If it is not prepared as part of the employee's official duties, it may be subject to copyright.

If "Government" is chosen, please do not choose a Creative Commons License. The work will be published with "Written work prepared by employees of the Federal Government as part of their official duties is, under the U.S. Copyright Act, a "work of the United States Government" for which copyright protection under Title 17 of the United States Code is not available. As such, copyright does not extend to the contributions of employees of the Federal Government."

NOTE: If author has selected Item 4, Section 3. on the Agreement will not apply to the Work.

Note: The work was funded by Amgen and we wish to may for Gold Open Access; please invoice the author for correspondence

SIGNATURE PAGE

The Corresponding Author acknowledges and agrees that the Corresponding Author is entering into, and has executed, the Agreement on behalf of the Corresponding Author and each other author named as contributing to the Article (each such author, an “Author”, and collectively, the “Authors”). The Corresponding Author represents and warrants that the Corresponding Author has obtained permission from each Author to enter into the Agreement on behalf of such Author and the Corresponding Author and each Author has read, understands, and has agreed to the terms of the Agreement, including, without limitation, the terms contained in the Agreement with respect to authorized reuse of the Article.

IN WITNESS WHEREOF, the Author has executed this License, effective as of the Effective Date.

PRINT NAME

SIGNATURE

Important Note: Once you electronically sign this form, you will not be able to make any additional changes to it.

To electronically sign this form, click the signature field above and provide the information requested in the dialog boxes.



[Click here to access/download](#)

**Copyright Transfer Agreement--REQUIRED from ALL
authors of submission at revision stage**
CTA Bryan.pdf





[Click here to access/download](#)

**Copyright Transfer Agreement--REQUIRED from ALL
authors of submission at revision stage**

CTA Cen.pdf





[Click here to access/download](#)

**Copyright Transfer Agreement--REQUIRED from ALL
authors of submission at revision stage**

CTA Dawn.pdf





[Click here to access/download](#)

**Copyright Transfer Agreement--REQUIRED from ALL
authors of submission at revision stage**
CTA Goadsby.pdf





[Click here to access/download](#)

**Copyright Transfer Agreement--REQUIRED from ALL
authors of submission at revision stage**

CTA Hantao.jpg





[Click here to access/download](#)

**Copyright Transfer Agreement--REQUIRED from ALL
authors of submission at revision stage
CTA Hong.pdf**



[Click here to access/download](#)

**Copyright Transfer Agreement--REQUIRED from ALL
authors of submission at revision stage**

CTA Jan.pdf





[Click here to access/download](#)

**Copyright Transfer Agreement--REQUIRED from ALL
authors of submission at revision stage**

CTA Judy.pdf





[Click here to access/download](#)

**Copyright Transfer Agreement--REQUIRED from ALL
authors of submission at revision stage**

CTA Licheng.pdf





[Click here to access/download](#)

**Copyright Transfer Agreement--REQUIRED from ALL
authors of submission at revision stage**

CTA Margarida.pdf





[Click here to access/download](#)

**Copyright Transfer Agreement--REQUIRED from ALL
authors of submission at revision stage**

CTA Ruoyuan.pdf





[Click here to access/download](#)

**Copyright Transfer Agreement--REQUIRED from ALL
authors of submission at revision stage**

CTA Silke.pdf





[Click here to access/download](#)

**Copyright Transfer Agreement--REQUIRED from ALL
authors of submission at revision stage**

CTA Simon.pdf





[Click here to access/download](#)

**Copyright Transfer Agreement--REQUIRED from ALL
authors of submission at revision stage**

CTA Sonya.pdf





[Click here to access/download](#)

**Copyright Transfer Agreement--REQUIRED from ALL
authors of submission at revision stage
CTA Weera.pdf**

Please wait...

If this message is not eventually replaced by the proper contents of the document, your PDF viewer may not be able to display this type of document.

You can upgrade to the latest version of Adobe Reader for Windows®, Mac, or Linux® by visiting http://www.adobe.com/go/reader_download.

For more assistance with Adobe Reader visit <http://www.adobe.com/go/acrreader>.

Windows is either a registered trademark or a trademark of Microsoft Corporation in the United States and/or other countries. Mac is a trademark of Apple Inc., registered in the United States and other countries. Linux is the registered trademark of Linus Torvalds in the U.S. and other countries.

Please wait...

If this message is not eventually replaced by the proper contents of the document, your PDF viewer may not be able to display this type of document.

You can upgrade to the latest version of Adobe Reader for Windows®, Mac, or Linux® by visiting <http://www.adobe.com/products/acrobat/readstep2.html>.

For more assistance with Adobe Reader visit <http://www.adobe.com/support/products/acrreader.html>.

Windows is either a registered trademark or a trademark of Microsoft Corporation in the United States and/or other countries. Mac is a trademark of Apple Inc., registered in the United States and other countries. Linux is the registered trademark of Linus Torvalds in the U.S. and other countries.

Please wait...

If this message is not eventually replaced by the proper contents of the document, your PDF viewer may not be able to display this type of document.

You can upgrade to the latest version of Adobe Reader for Windows®, Mac, or Linux® by visiting http://www.adobe.com/go/reader_download.

For more assistance with Adobe Reader visit <http://www.adobe.com/go/acrreader>.

Windows is either a registered trademark or a trademark of Microsoft Corporation in the United States and/or other countries. Mac is a trademark of Apple Inc., registered in the United States and other countries. Linux is the registered trademark of Linus Torvalds in the U.S. and other countries.

Please wait...

If this message is not eventually replaced by the proper contents of the document, your PDF viewer may not be able to display this type of document.

You can upgrade to the latest version of Adobe Reader for Windows®, Mac, or Linux® by visiting <http://www.adobe.com/products/acrobat/readstep2.html>.

For more assistance with Adobe Reader visit <http://www.adobe.com/support/products/acrreader.html>.

Windows is either a registered trademark or a trademark of Microsoft Corporation in the United States and/or other countries. Mac is a trademark of Apple Inc., registered in the United States and other countries. Linux is the registered trademark of Linus Torvalds in the U.S. and other countries.

ICMJE Form for Disclosure of Potential Conflicts of Interest

Instructions

The purpose of this form is to provide readers of your manuscript with information about your other interests that could influence how they receive and understand your work. The form is designed to be completed electronically and stored electronically. It contains programming that allows appropriate data display. Each author should submit a separate form and is responsible for the accuracy and completeness of the submitted information. The form is in six parts.

1. Identifying information.

2. The work under consideration for publication.

This section asks for information about the work that you have submitted for publication. The time frame for this reporting is that of the work itself, from the initial conception and planning to the present. The requested information is about resources that you received, either directly or indirectly (via your institution), to enable you to complete the work. Checking "No" means that you did the work without receiving any financial support from any third party -- that is, the work was supported by funds from the same institution that pays your salary and that institution did not receive third-party funds with which to pay you. If you or your institution received funds from a third party to support the work, such as a government granting agency, charitable foundation or commercial sponsor, check "Yes".

3. Relevant financial activities outside the submitted work.

This section asks about your financial relationships with entities in the bio-medical arena that could be perceived to influence, or that give the appearance of potentially influencing, what you wrote in the submitted work. You should disclose interactions with ANY entity that could be considered broadly relevant to the work. For example, if your article is about testing an epidermal growth factor receptor (EGFR) antagonist in lung cancer, you should report all associations with entities pursuing diagnostic or therapeutic strategies in cancer in general, not just in the area of EGFR or lung cancer.

Report all sources of revenue paid (or promised to be paid) directly to you or your institution on your behalf over the 36 months prior to submission of the work. This should include all monies from sources with relevance to the submitted work, not just monies from the entity that sponsored the research. Please note that your interactions with the work's sponsor that are outside the submitted work should also be listed here. If there is any question, it is usually better to disclose a relationship than not to do so.

For grants you have received for work outside the submitted work, you should disclose support ONLY from entities that could be perceived to be affected financially by the published work, such as drug companies, or foundations supported by entities that could be perceived to have a financial stake in the outcome. Public funding sources, such as government agencies, charitable foundations or academic institutions, need not be disclosed. For example, if a government agency sponsored a study in which you have been involved and drugs were provided by a pharmaceutical company, you need only list the pharmaceutical company.

4. Intellectual Property.

This section asks about patents and copyrights, whether pending, issued, licensed and/or receiving royalties.

5. Relationships not covered above.

Use this section to report other relationships or activities that readers could perceive to have influenced, or that give the appearance of potentially influencing, what you wrote in the submitted work.

Definitions.

Entity: government agency, foundation, commercial sponsor, academic institution, etc.

Grant: A grant from an entity, generally [but not always] paid to your organization

Personal Fees: Monies paid to you for services rendered, generally honoraria, royalties, or fees for consulting, lectures, speakers bureaus, expert testimony, employment, or other affiliations

Non-Financial Support: Examples include drugs/equipment supplied by the entity, travel paid by the entity, writing assistance, administrative support, etc.

Other: Anything not covered under the previous three boxes

Pending: The patent has been filed but not issued

Issued: The patent has been issued by the agency

Licensed: The patent has been licensed to an entity, whether earning royalties or not

Royalties: Funds are coming in to you or your institution due to your patent

ICMJE Form for Disclosure of Potential Conflicts of Interest

Section 1. Identifying Information

1. Given Name (First Name)

2. Surname (Last Name)

3. Date

4. Are you the corresponding author?

☐ Yes ☒ No

Corresponding Author's Name

Peter J. Goadsby

5. Manuscript Title

PAC1 receptor blockade reduces central nociceptive activity- new approach for primary headache?

6. Manuscript Identifying Number (if you know it)

PAIN-D-19-01173

Section 2. The Work Under Consideration for Publication

Did you or your institution **at any time** receive payment or services from a third party (government, commercial, private foundation, etc.) for any aspect of the submitted work (including but not limited to grants, data monitoring board, study design, manuscript preparation, statistical analysis, etc.)?

Are there any relevant conflicts of interest? ☐ Yes ☐ No

Section 3. Relevant financial activities outside the submitted work.

Place a check in the appropriate boxes in the table to indicate whether you have financial relationships (regardless of amount of compensation) with entities as described in the instructions. Use one line for each entity; add as many lines as you need by clicking the "Add +" box. You should report relationships that were **present during the 36 months prior to publication**.

Are there any relevant conflicts of interest? ☐ Yes ☐ No

Section 4. Intellectual Property -- Patents & Copyrights

Do you have any patents, whether planned, pending or issued, broadly relevant to the work? ☐ Yes ☐ No

ICMJE Form for Disclosure of Potential Conflicts of Interest

Section 5. Relationships not covered above

Are there other relationships or activities that readers could perceive to have influenced, or that give the appearance of potentially influencing, what you wrote in the submitted work?

- ☐ Yes, the following relationships/conditions/circumstances are present (explain below):
- ☐ No other relationships/conditions/circumstances that present a potential conflict of interest

At the time of manuscript acceptance, journals will ask authors to confirm and, if necessary, update their disclosure statements. On occasion, journals may ask authors to disclose further information about reported relationships.

Section 6. Disclosure Statement

Based on the above disclosures, this form will automatically generate a disclosure statement, which will appear in the box below.

Evaluation and Feedback

Please visit <http://www.icmje.org/cgi-bin/feedback> to provide feedback on your experience with completing this form.

Please wait...

If this message is not eventually replaced by the proper contents of the document, your PDF viewer may not be able to display this type of document.

You can upgrade to the latest version of Adobe Reader for Windows®, Mac, or Linux® by visiting http://www.adobe.com/go/reader_download.

For more assistance with Adobe Reader visit <http://www.adobe.com/go/acrreader>.

Windows is either a registered trademark or a trademark of Microsoft Corporation in the United States and/or other countries. Mac is a trademark of Apple Inc., registered in the United States and other countries. Linux is the registered trademark of Linus Torvalds in the U.S. and other countries.

Please wait...

If this message is not eventually replaced by the proper contents of the document, your PDF viewer may not be able to display this type of document.

You can upgrade to the latest version of Adobe Reader for Windows®, Mac, or Linux® by visiting <http://www.adobe.com/products/acrobat/readstep2.html>.

For more assistance with Adobe Reader visit <http://www.adobe.com/support/products/acrreader.html>.

Windows is either a registered trademark or a trademark of Microsoft Corporation in the United States and/or other countries. Mac is a trademark of Apple Inc., registered in the United States and other countries. Linux is the registered trademark of Linus Torvalds in the U.S. and other countries.

Please wait...

If this message is not eventually replaced by the proper contents of the document, your PDF viewer may not be able to display this type of document.

You can upgrade to the latest version of Adobe Reader for Windows®, Mac, or Linux® by visiting http://www.adobe.com/go/reader_download.

For more assistance with Adobe Reader visit <http://www.adobe.com/go/acrreader>.

Windows is either a registered trademark or a trademark of Microsoft Corporation in the United States and/or other countries. Mac is a trademark of Apple Inc., registered in the United States and other countries. Linux is the registered trademark of Linus Torvalds in the U.S. and other countries.

Please wait...

If this message is not eventually replaced by the proper contents of the document, your PDF viewer may not be able to display this type of document.

You can upgrade to the latest version of Adobe Reader for Windows®, Mac, or Linux® by visiting http://www.adobe.com/go/reader_download.

For more assistance with Adobe Reader visit <http://www.adobe.com/go/acrreader>.

Windows is either a registered trademark or a trademark of Microsoft Corporation in the United States and/or other countries. Mac is a trademark of Apple Inc., registered in the United States and other countries. Linux is the registered trademark of Linus Torvalds in the U.S. and other countries.

Please wait...

If this message is not eventually replaced by the proper contents of the document, your PDF viewer may not be able to display this type of document.

You can upgrade to the latest version of Adobe Reader for Windows®, Mac, or Linux® by visiting http://www.adobe.com/go/reader_download.

For more assistance with Adobe Reader visit <http://www.adobe.com/go/acrreader>.

Windows is either a registered trademark or a trademark of Microsoft Corporation in the United States and/or other countries. Mac is a trademark of Apple Inc., registered in the United States and other countries. Linux is the registered trademark of Linus Torvalds in the U.S. and other countries.

Please wait...

If this message is not eventually replaced by the proper contents of the document, your PDF viewer may not be able to display this type of document.

You can upgrade to the latest version of Adobe Reader for Windows®, Mac, or Linux® by visiting http://www.adobe.com/go/reader_download.

For more assistance with Adobe Reader visit <http://www.adobe.com/go/acrreader>.

Windows is either a registered trademark or a trademark of Microsoft Corporation in the United States and/or other countries. Mac is a trademark of Apple Inc., registered in the United States and other countries. Linux is the registered trademark of Linus Torvalds in the U.S. and other countries.

Please wait...

If this message is not eventually replaced by the proper contents of the document, your PDF viewer may not be able to display this type of document.

You can upgrade to the latest version of Adobe Reader for Windows®, Mac, or Linux® by visiting http://www.adobe.com/go/reader_download.

For more assistance with Adobe Reader visit <http://www.adobe.com/go/acrreader>.

Windows is either a registered trademark or a trademark of Microsoft Corporation in the United States and/or other countries. Mac is a trademark of Apple Inc., registered in the United States and other countries. Linux is the registered trademark of Linus Torvalds in the U.S. and other countries.

Please wait...

If this message is not eventually replaced by the proper contents of the document, your PDF viewer may not be able to display this type of document.

You can upgrade to the latest version of Adobe Reader for Windows®, Mac, or Linux® by visiting http://www.adobe.com/go/reader_download.

For more assistance with Adobe Reader visit <http://www.adobe.com/go/acrreader>.

Windows is either a registered trademark or a trademark of Microsoft Corporation in the United States and/or other countries. Mac is a trademark of Apple Inc., registered in the United States and other countries. Linux is the registered trademark of Linus Torvalds in the U.S. and other countries.

Please wait...

If this message is not eventually replaced by the proper contents of the document, your PDF viewer may not be able to display this type of document.

You can upgrade to the latest version of Adobe Reader for Windows®, Mac, or Linux® by visiting <http://www.adobe.com/products/acrobat/readstep2.html>.

For more assistance with Adobe Reader visit <http://www.adobe.com/support/products/acrreader.html>.

Windows is either a registered trademark or a trademark of Microsoft Corporation in the United States and/or other countries. Mac is a trademark of Apple Inc., registered in the United States and other countries. Linux is the registered trademark of Linus Torvalds in the U.S. and other countries.

Please wait...

If this message is not eventually replaced by the proper contents of the document, your PDF viewer may not be able to display this type of document.

You can upgrade to the latest version of Adobe Reader for Windows®, Mac, or Linux® by visiting <http://www.adobe.com/products/acrobat/readstep2.html>.

For more assistance with Adobe Reader visit <http://www.adobe.com/support/products/acrreader.html>.

Windows is either a registered trademark or a trademark of Microsoft Corporation in the United States and/or other countries. Mac is a trademark of Apple Inc., registered in the United States and other countries. Linux is the registered trademark of Linus Torvalds in the U.S. and other countries.